



Holomorphic normal form of nonlinear perturbations of nilpotent vector fields

Laurent Stolovitch, Freek Verstringe

► To cite this version:

Laurent Stolovitch, Freek Verstringe. Holomorphic normal form of nonlinear perturbations of nilpotent vector fields. *Regular and Chaotic Dynamics*, 2016, 21, pp.410-436. 10.1134/S1560354716040031 . hal-01336282

HAL Id: hal-01336282

<https://hal.science/hal-01336282>

Submitted on 22 Jun 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Holomorphic normal form of nonlinear perturbations of nilpotent vector fields

Laurent Stolovitch ^{*}and Freek Verstringe [†]

June 22, 2016

Abstract

We consider germs of holomorphic vector fields at a fixed point having a nilpotent linear part at that point, in dimension $n \geq 3$. Based on Belitskii's work, we know that such a vector field is formally conjugate to a (formal) normal form. We give a condition on that normal form which ensure that the normalizing transformation is holomorphic at the fixed point. We shall show that this sufficient condition is a *nilpotent version* of Bruno's condition (A). In dimension 2, no condition is required since, according to Stróżyńska-Żoładek, each such germ is holomorphically conjugate to a Takens normal form. Our proof is based on Newton's method and $\mathfrak{sl}_2(\mathbb{C})$ -representations.

1 Introduction

In this article, we consider germs of holomorphic vector fields in a neighborhood of a fixed point in \mathbb{C}^n , $n \geq 2$. We are interested in the local classification under the action of the group of germs of biholomorphisms preserving the fixed point, which we may assume to be the origin. In the sequel, “germ” of vector field or map refers to a holomorphic germ at the origin. The idea is to simplify, by a change of coordinates, the system of differential equations in order to better understand its dynamics. An elementary instance of such a problem is the following : Let A be a $n \times n$ -matrix with complex coefficients. In order to understand the orbits $\{A^k z\}_{k \in \mathbb{Z}}$, z near the fixed point 0 of A , it is much easier to transform A into its Jordan normal form J by a change of coordinates P , $A = PJP^{-1}$. Then, one studies $\{J^k y\}_{k \in \mathbb{Z}}$, $Py = z$ and pulls back the information to the original problem. It was the idea of Poincaré to

^{*}CNRS, Laboratoire J.-A. Dieudonné U.M.R. 6621, Université de Nice - Sophia Antipolis, Parc Valrose 06108 Nice Cedex 02, France. email : `stolo@unice.fr` Research of L. Stolovitch was supported by ANR grant “ANR-10-BLAN 0102” for the project DynPDE

[†]Royal Observatory of Belgium, Ringlaan 3, 1180 Brussels, Belgium. email : `freek.verstringe@oma.be`

develop this point of view for vector fields. The difficulty is that the Lie algebra of germs of vector fields is infinite dimensional. The linear transformations are replaced by germs of diffeomorphisms fixing the origin. The special representant of the orbit of the action that plays the role of the Jordan normal form is called a normal form. A precise definition is given below. This idea has been extensively developed by Arnold and his school, Bruno, Moser, Ecalle, Martinet-Ramis, Yoccoz, ... in the case where the linear part at the fixed point is a semi-simple matrix. The issue is that, although conjugacy to a formal normal form can always be obtained by a suitable formal transformation, it may not be possible to reach a normal form by a holomorphic transformation at the fixed point.

In this article, we investigate the normal form problem, in dimension $n \geq 3$, for holomorphic vector fields with a nilpotent linear part at the fixed point (assumed to be the origin). We give a sufficient condition that ensures that a germ of holomorphic vector field can be holomorphically conjugated to a normal form.

1.1 Formal normal forms

Following G. Belitskii [Bel79, Bel82], E. Lombardi and L. Stolovitch defined in [LS10], a notion of normal form of higher order perturbations of a given quasi-homogeneous vector field S and the associated notion of “generalized resonances” (these are obstructions to find a formal power series transformation conjugating the perturbation of S back to S itself). Here, we shall first focus on the case where S is linear, and then on the case where S is nilpotent.

Let \mathcal{H}_k be the space of homogeneous vector fields of degree k . For each natural number $k \geq 2$, we consider the “cohomological operator” $d_0 := [S, \cdot] : \mathcal{H}_k \rightarrow \mathcal{H}_k$ associated to S . Here, $[S, \cdot]$ denotes the Lie bracket with S . We define a space of normal forms of degree k to be a supplementary space \mathcal{C}_k to $\text{Im} d_0|_{\mathcal{H}_k}$. We can show that there exists a formal transformation that conjugates each perturbation of S of order ≥ 2 , $X := S + R_{\geq 2}$, to a formal vector field $NF := S + \sum_{k \geq 2} v_k$ where $v_k \in \mathcal{C}_k$ for each k . We call NF a normal form (of “style” $\hat{\mathcal{C}} = \oplus \mathcal{C}_j$, see [Mur04]). Indeed, by induction on the degree $k \geq 2$, assume that X is normalized up to order $k-1$, that is $X = NF^{k-1} + R_{\geq k}$ and $NF^{k-1} \in \oplus_{j=2}^{k-1} \mathcal{C}_j$. Let $\pi_{\text{Im } d_0}$ denote the projection onto the range of d_0 along \mathcal{C}_k . Let us choose $U_k \in \mathcal{H}_k$ to solve the **cohomological equation** $d_0(U_k) = -\pi_{\text{Im } d_0} R_k$, where R_k denotes the homogeneous polynomial of degree k of $R_{\geq k}$. Then $(I - \pi_{\text{Im } d_0})R_k \in \mathcal{C}_k$ and we have,

$$(\text{id} + U_k)_*^{-1} X = NF_{k-1} + [S, U_k] + R_k + h.o.t. = NF_k + R_{>k},$$

where $NF_k := NF_{k-1} + (I - \pi_{\text{Im } d_0})R_k$. Here, $\Phi_* X := D\Phi(\Phi^{-1})X(\Phi^{-1})$ denotes the conjugacy of X by the diffeomorphism Φ . It is classical that a normal form is not uniquely defined since one can add any $U_0 \in \text{Ker } d_0|_{\mathcal{H}_k}$ to the solution U_k of $d_0(U_k) = -\pi_{\text{Im } d_0} R_k$. Therefore, defining for any $k \geq 2$, a supplementary space \tilde{V}_k to $\text{Ker } d_0|_{\mathcal{H}_k}$ in \mathcal{H}_k , we can restrict the solution U_k to belong to \tilde{V}_k , for all k , and refer

to the normal form so obtained as *the* normal form of X . The corresponding formal transformation $\cdots \circ (\text{id} + U_k) \circ \cdots \circ (\text{id} + U_1)$ is called *the associated normalizing transformation*.

Of course, we are interested in the case where these supplementary spaces can be computed effectively. One way to do so is as follows : \mathcal{H}_k is provided with an Hermitian scalar product (see (3)). Let us define \mathcal{C}_k (resp. \tilde{V}_k) to be the orthogonal complement to $\text{Im}(d_{0|\mathcal{H}_k})$ (resp. to be equal to $\text{Im}(d_{0|\mathcal{H}_k}^*)$), that is $\mathcal{H}_k = \text{Im}(d_{0|\mathcal{H}_k}) \oplus^\perp \mathcal{C}_k$. The space \mathcal{C}_k is the kernel of the adjoint $d_{0|\mathcal{H}_k}^*$ of $d_{0|\mathcal{H}_k}$ with respect to the Hermitian scalar product.

When S is a semi-simple linear vector field with eigenvalues $\lambda_1, \dots, \lambda_n$, the space of normal forms is defined in terms of resonance relations $\sum_j q_j \lambda_j = \lambda_i$: the space \mathcal{C}_k generated by monomial vector fields $x^Q \partial_{x_i}$ with $Q = (q_1, \dots, q_n) \in \mathbb{N}^n$, $|Q| = q_1 + \cdots + q_n = k \geq 2$, $1 \leq i \leq n$ which satisfies a resonance relation. We can also choose $\tilde{V}_k = \mathcal{C}_k$ and obtain the classical Poincaré-Dulac normal form [Arn80].

When S is nilpotent, the normal form spaces are more diverse and there are different ways of defining normal forms. In dimension $n = 2$, F. Takens [Tak73] has shown that the space of normal forms associated to $S = y \partial_x$, \mathcal{C}_k , can be chosen to be the vector space generated by $x^k \partial_x$ and $x^k \partial_y$, $k \geq 2$. This has been generalized recently by E. Stróżyńska and H. Żoładek [SZ08] to the case $n \geq 2$.

There is another classical way to define normal forms of perturbations of nilpotent linear vector fields, by considering an $\mathfrak{sl}(2, \mathbb{C})$ -triple associated to S and their representations (see [CS86, Mur03]). First of all, it is classical (Jacobson-Morozov theorem)[Ser87, Bou90] that there exists linear vector fields $N := S$, M and H of \mathbb{C}^n such that $[N, M] = H$, $[H, N] = 2N$, $[H, M] = -2M$. A representation of $\mathfrak{sl}(2, \mathbb{C})$ in a finite dimensional vector space V , is a triple of endomorphisms X, Y, Z such that

$$[X, Y] = Z, \quad [Z, X] = 2X, \quad [Z, Y] = -2Y.$$

Here, $[X, Y]$ denotes the bracket of endomorphism $XY - YX$. Such a family $\{X, Y, Z\}$ is called an $\mathfrak{sl}(2, \mathbb{C})$ -triple. It is classical that $V = \text{Im} X \oplus \text{Ker} Y$. We apply this to $X = [N, \cdot]$, $Y = [M, \cdot]$ acting on the space of homogeneous polynomial vector fields : The supplementary space to the image of d_0 is then defined to be the kernel of $\text{ad}_M := [M, \cdot]$ restricted to the space of homogeneous polynomial vector fields.

There exist also more involved constructions of formal normal forms dealing with uniqueness problems such as in [KOW96] for perturbation of (quasi-)homogeneous vector fields or in [San03, GY12], based on spectral sequences.

1.2 Analytic normal form problem

As we have seen, for each higher order perturbation of S , there exists a formal transformation (i.e a formal change of coordinates) to a formal normal form. We are interested in the situation where not only the perturbation is analytic in a neighbor-

hood of the origin but also the normalizing transformation to its normal form (which is thus analytic too).

Hence we are led to solve and estimate iteratively the solution U_k of the “cohomological equation” $d_0(U_k) = F_k$ for a given polynomial F_k of degree k in the range of d_0 .

If S is semi-simple, say $S = \sum_{i=1}^n \lambda_i x_i \frac{\partial}{\partial x_i}$, this leads to the so-called *small divisor problem* [SM71, Mos73, Arn80] : the norm of U_k is bounded by the norm of F_k divided by smallest of the non-zero numbers of the form $|q_1 \lambda_1 + \dots + q_n \lambda_n - \lambda_j|$, $Q = (q_1, \dots, q_n) \in \mathbb{N}^n$, $|Q| := q_1 + \dots + q_n = k$. These numbers may accumulate to the origin when the degree k tends to $+\infty$ causing a blow up of the normalizing transformation. These numbers are called “small divisors” (although they may not be small). The faster these numbers accumulate to zero, the smaller the radius of convergence of the formal transformation is. For instance, C. L. Siegel proved [Sie42], that if there exists $\tau \geq 0$ such that $|q_1 \lambda_1 + \dots + q_n \lambda_n - \lambda_j| > \frac{C}{|Q|^\tau}$ for all $Q \in \mathbb{N}^n$, $|Q| \geq 2$, then any higher order analytic perturbation of S is analytically linearizable. This condition has been weakened to the so-called (ω) -condition by Bruno [Bru72] (we shall not recall here its definition since we shall not use it). Nevertheless, the control of these small divisors is not sufficient to ensure the analyticity of the normalizing transformation to a (non-linear) normal form. Indeed, there are well known situation for which, although the small divisors are bounded away from the origin, the normalizing transformation is anyway divergent at the origin [Mar80]. Finally, A.D. Bruno [Bru72] found a necessary and sufficient condition on the formal normal form - condition (A)- that ensures that if the semi-simple linear part satisfies the diophantine condition (ω) , then any analytic perturbation of S that has a formal normal form satisfying (A) can be analytically normalized. This is a generalization of H. Rüssmann’s Hamiltonian version [Rüs67]. These works have been generalized to several commuting vector fields in several directions [Ito89, Sto00, Sto05, Zun05, Zun02, Ito09] in the spirit of J. Vey [Vey79, Vey78].

In [LS10], a broader notion of “small divisors” associated to a linear vector field S is defined. They are defined as the square roots of the nonzero eigenvalues of the “box operator” $\square_{\mathcal{H}_k} := d_{0|\mathcal{H}_k} d_{0|\mathcal{H}_k}^*$, for all $k > 2$. When S is a nilpotent linear vector field in dimension two and three, G. Iooss and E. Lombardi [IL05] computed the corresponding “generalized resonances” as well as the small divisors. Recently, P. Bonckaert and F. Verstringe succeeded in [BV12], to estimate the generalized small divisors for (almost) any nilpotent linear part. It is there shown that they are always bounded away from the origin.

One of the remaining questions was whether a Takens normal form (in dimension 2) could be obtained systematically by a convergent transformation. It took almost forty years to be answered. In dimension 2, after an attempt to solve this question by X. Gong [Gon95], finally E. Stróżyńska and H. Żoładek [SZ02] showed that any holomorphic nonlinear perturbation of $y \frac{\partial}{\partial x}$ has a germ of biholomorphism that conjugate it to a Takens normal form : $(y + f(x, y)) \frac{\partial}{\partial x} + g(x, y) \frac{\partial}{\partial y}$ is analytically conjugate to

$(y + k(x))\frac{\partial}{\partial x} + l(x)\frac{\partial}{\partial y}$; here f, g, k, l are analytic germs of order ≥ 2 at the origin. A more geometric proof was given in [Lor06]. It was shown also by E. Stróżyńska and H. Żoładek that the equivalent theorem in a higher dimensional setting is false [SŻ11, SŻ08].

For the similar problem with the “scalar product normal form”, we emphasize, that in dimension 2, the “small divisors” tend to infinity with the degree of homogeneity. This phenomenon has a regularizing effect on the solution of the conjugacy equation to a normal form. This corresponds, somehow, to the situation of the Poincaré domain for non-zero semi-simple linear part, where no requirements are needed on the perturbation for it to have an analytic transformation to a normal form.

In higher dimension an example in [IL05] shows that the “small divisors” may not tend to infinity with the degree of homogeneity : they may accumulate finite numbers as well as infinity. The higher dimensional situation for the nilpotent case corresponds somehow to the “Siegel domain” : one needs to impose conditions on the formal normal form in order to obtain the convergence of such a transformation.

The aim of this article is to investigate, in dimension $n \geq 3$, the holomorphic conjugacy to a normal form problem for an higher order analytic perturbation of nilpotent linear vector field. We shall define a nilpotent version of Bruno’s condition (A). We shall state and prove a nilpotent version of the “sufficiency part” of Bruno’s theorem : in dimension ≥ 3 , if an analytic higher perturbation of a linear nilpotent N has its formal normal form that satisfies this condition, then there exists an analytic transformation to its normal form.

1.3 Main result

In this section we formulate our main result using the notations introduced in the introduction. We first introduce some extra terminology. Let S be a linear vector field. We denote by $\hat{\mathcal{O}}_n^S$ the ring of formal first integrals of S , that is the set of formal power series \hat{f} for which the Lie derivative $\mathcal{L}_S(\hat{f}) = 0$ vanishes. A nilpotent matrix is said to be *regular* if its Jordan normal form does not contain zero blocks. In other words, a nilpotent endomorphism is regular if there is no invariant subspace upon which the operator acts as a zero operator. We shall denote by $R_{\geq 2}$ a germ of holomorphic vector field (or function, map depending on the context) vanishing at order ≥ 2 at the origin.

Theorem 1.1. *Let N be a regular nilpotent linear vector field of \mathbb{C}^n and assume that $\{N, N^*, [N, N^*]\}$ is a $\mathfrak{sl}(2, \mathbb{C})$ -triple. Let $V = N + R_{\geq 2}$ be a germ of holomorphic vector field at the origin of \mathbb{C}^n . Assume that its formal normal form (as defined by the iteration procedure described in the introduction) has the form*

$$V = N + fN^* \quad \text{where} \quad f \in \hat{\mathcal{O}}_n^N \cap \hat{\mathcal{O}}_n^{N^*}. \quad (1)$$

Then, its associated formal diffeomorphism converges at the origin.

This is the main result of our article and should be understood as a “**complete integrability**” **theorem** along the same lines as (and in fact a continuation of) results by A.D. Bruno [Bru72] (see also [Mar80, Zun02]), and by the first author [Sto00, Sto05]. Indeed, in these articles, the authors consider a (germ of) vector field (or commuting families of vector fields) $V = S + R_{\geq 2}$ that are holomorphic nonlinear perturbation of a semi-simple linear vector field $S = \sum_{i=1}^n \lambda_i x_i \frac{\partial}{\partial x_i}$. The main (easiest) theorem of Bruno can be stated as follows : Assume that the normal form of V is of the form $S + fS + g\bar{S}$ where $\bar{S} := \sum_{i=1}^n \bar{\lambda}_i x_i \frac{\partial}{\partial x_i}$ and $f, g \in \hat{\mathcal{O}}_n^S$, $f(0) = g(0) = 0$ (this is Bruno’s “condition (A)”). If S satisfies Bruno’s small divisors condition (ω), then its associated formal biholomorphism is convergent at the origin and it conjugates V to a (convergent) normal form.

Theorem 1.1 can be seen as a **nilpotent version of Bruno’s theorem**. Indeed, \bar{S} plays the role of N^* and we have $\hat{\mathcal{O}}_n^S = \hat{\mathcal{O}}_n^S \cap \hat{\mathcal{O}}_n^{\bar{S}}$. In our normal form (1), there is no equivalent term to fS since the natural one would be fN which is not a normal form w.r.t. to N , i.e. $fN \notin \text{Ker } \text{ad}_{N^*}$.

Remark 1.2. *One of the key points of [BV12] is that, if the linear nilpotent vector field S is regular, then there exists a linear change of coordinates L of \mathbb{C}^n such that $N := L_*S$, N^* , the vector field defined as the adjoint of the differential operator acting on polynomials of N w.r.t. the scalar product used above, and $H = [N, N^*]$ define an $\mathfrak{sl}(2, \mathbb{C})$ -triple. Moreover, $\{\text{ad}_N, \text{ad}_{N^*}, \text{ad}_N \text{ad}_{N^*} - \text{ad}_{N^*} \text{ad}_N\}$ is also an $\mathfrak{sl}(2, \mathbb{C})$ -triple. In the sequel, we shall assume that the linear change of coordinates L has been applied and hence, that both $\{N, N^*, [N, N^*]\}$ and $\{\text{ad}_N, \text{ad}_{N^*}, [\text{ad}_N, \text{ad}_{N^*}]\}$ are $\mathfrak{sl}(2, \mathbb{C})$ -triples.*

Remark 1.3. *If $V = N + R_{\geq 2}$ is a germ of holomorphic vector field at the origin of \mathbb{C}^n that can be embedded into an analytic $\mathfrak{sl}(2, \mathbb{C})$ -triple, that is if there are germs $\tilde{V} = N^* + \text{h.o.t.}$, $V' = H + \text{h.o.t.}$ of holomorphic vector fields such that $\{V, \tilde{V}, V'\}$ is (Lie-)isomorphic to $\mathfrak{sl}(2, \mathbb{C})$, then V, \tilde{V} and V' are simultaneously and holomorphically linearizable. Indeed, at the formal level, this follows from [Her68] while the analytic result follows from [GS68, Kus67]. It should be emphasized that this result does not hold in the smooth category [GS68, CG97].*

Remark 1.4. *If $V = N + R_{\geq 2}$, a germ of holomorphic vector field at the origin of \mathbb{C}^n , can be embedded into a formal $\mathfrak{sl}(2, \mathbb{C})$ -triple, that is if there are formal vector fields $\tilde{V} = N^* + \text{h.o.t.}$, $V' = H + \text{h.o.t.}$ such that $\{V, \tilde{V}, V'\}$ is (Lie-)isomorphic to $\mathfrak{sl}(2, \mathbb{C})$, then V, \tilde{V} and V' are simultaneously formally linearizable. According to [BV12][p.2223], the “small divisors” are bounded away from the origin. Then, according to [LS10][theorem 5.8], V is analytically linearizable.*

There are other results concerning the problem of classification of perturbations not of nilpotent vector fields but rather of quasihomogenous vector fields with a nilpotent linear part at the origin. They are all in dimension 2 (see for instance [MC88, CDS04]) and are not immediately concerned with holomorphic conjugacy (in a neighborhood) at the origin to a normal form.

1.4 Geometric interpretation

According to Weitzenböck's theorem [Wei32], the ring of formal first integrals $\widehat{\mathcal{O}}_n^N$ of any nilpotent linear vector field N is finitely generated over \mathbb{C} (see [Now94][theorem 6.2.1] for this formulation).

The ring of common formal first integrals $\widehat{\mathcal{O}}_n^N \cap \widehat{\mathcal{O}}_n^{N^*}$ is hence finitely generated over \mathbb{C} . Let P_1, \dots, P_r be a set of generating polynomials. If they are algebraically dependent, then there exists polynomials Q_1, \dots, Q_l on \mathbb{C}^r such that $Q_i(P_1, \dots, P_r) = 0$ for all i . Let us consider

$$\mathcal{C} = \{z \in \mathbb{C}^r \mid Q_i(z) = 0, i = 1, \dots, l\}$$

the zero locus of these polynomials. If the P_j 's are algebraically independent, then $\mathcal{C} = \mathbb{C}^r$. Consider the map

$$\begin{aligned} \pi : \mathbb{C}^n &\rightarrow (\mathcal{C}, 0), \\ x &\mapsto (P_1(x), \dots, P_r(x)). \end{aligned}$$

We apply our main result and assume that the vector field V has been conjugated into a normal form by a germ of holomorphic transformation at the origin. In these holomorphic coordinates, we have

$$V = N + fN^*$$

where $f \in \mathcal{O}_n^N \cap \mathcal{O}_n^{N^*}$ is a germ of holomorphic function. Let $b \in (\mathcal{C}, 0)$ be a point in the image of π . Then,

- (a) **V is tangent to the level set of $\Sigma_b := \pi^{-1}(b)$** , intersected with a neighborhood of the origin. Indeed, both N and N^* are tangent to this level set and the function f is constant on this level set.
- (b) the restriction of the vector field V , $V|_{\Sigma_b}$, **is the restriction to Σ_b of a linear vector field**.
- (c) the restriction $V|_{\Sigma_b}$, **belongs to the restriction an $\mathfrak{sl}(2, \mathbb{C})$ -action**¹, namely $\{N|_{\Sigma_b}, N^*|_{\Sigma_b}, H|_{\Sigma_b}\}$ (compare with remarks 1.3 and 1.4).
- (d) Moreover **this holds for every fiber** (in a neighborhood of the origin in \mathbb{C}^n) within a neighborhood of the origin in $\mathcal{C} \cap \text{Im } \pi$.

As a by-product, we also obtain that the zero fiber of π is invariant, the restriction to which V equals to the restriction of the linear nilpotent field N . As illustrated in the next example, this last result could have obtained under a much weaker assumption using [LS10][theorem 5.6] combined with the “no-small-divisors” statement of [BV12].

¹We thank Nguyen Tien Zung for this remark

Example 1.1. Consider the three dimensional linear vector field :

$$N := y \frac{\partial}{\partial x} + z \frac{\partial}{\partial y}.$$

We have

$$N^* := x \frac{\partial}{\partial y} + y \frac{\partial}{\partial z}.$$

Let us set

$$h = xz - \frac{y^2}{2}.$$

It follows that, $N(h) = 0 = N^*(h)$ and $\widehat{\mathcal{O}}_3^N \cap \widehat{\mathcal{O}}_3^{N^*} = \mathbb{C}[[h]]$. A formal normal form of a nonlinear perturbation of N is a formal vector field of the form

$$\begin{aligned} \dot{x} &= y + xP_1(x, h) \\ \dot{y} &= z + yP_1(x, h) + xP_2(x, h) \\ \dot{z} &= zP_1(x, h) + yP_2(x, h) + P_3(x, h) \end{aligned} \tag{2}$$

where the P_i 's are formal power series [IL05].

We apply our main result : Let $V = N + R_{\geq 2}$ be a germ of holomorphic vector field that is a nonlinear perturbation of N . Assume that the normal form of V is of the form :

$$\hat{\Phi}_* V = N + \hat{f} \left(xz - \frac{y^2}{2} \right) N^*$$

where \hat{f} is a formal power series of one variable (that is $P_1 = P_3 = 0$, $P_2(x, h) = \hat{f}(h)$). Then, $\hat{\Phi}$ defines a germ of holomorphic change of variables Φ at the origin of \mathbb{C}^3 and \hat{f} defines a germ of holomorphic function f at $0 \in \mathbb{C}$ such that

$$\Phi_* V = N + f \left(xz - \frac{y^2}{2} \right) N^*.$$

In these holomorphic coordinates, the vector field is tangent to each fiber

$$\Sigma_b = \left\{ (x, y, z) \in (\mathbb{C}^3, 0) \mid xz - \frac{y^2}{2} = b \right\}$$

in a neighborhood of the origin, b being sufficiently small. In particular, in these new holomorphic coordinates, the vector field is tangent to Σ_0 and its restriction to it is equal to (the restriction of) N . We emphasize that the last point can be achieved under a much weaker assumption : Assume that the formal normal form is such that $P_i(x, 0) = 0$, $i = 1, 2, 3$. Then, according to [LS10][theorem 5.6] applied to $\mathcal{I} = (h)$, there are holomorphic coordinates for which Σ_0 is an invariant set of the initial vector field X . Furthermore, its restriction to Σ_0 is equal to N .

1.5 Sketch of the proof

The proof of the main theorem is done using a Newton scheme. We assume that $V = NF_m + R_{\geq m+1}$ is normalized up to order $m = 2^k$. By assumption, the partial normal form NF_m is of the form $N + f_m N^*$ where $f_m \in \widehat{\mathcal{O}}_n^N \cap \widehat{\mathcal{O}}_n^{N^*}$ is a polynomial of degree $\leq m-1$. There is a unique polynomial vector field U of the form $U = d_0^*(V)$, of order $\geq m+1$ and of degree $\leq 2m$ such that the diffeomorphism $(\text{id} + U)^{-1}$ normalizes V up to order $2m$. The vector field U solves an equation of the form :

$$J^{2m}([NF_m, U]) = B$$

for some known B , in the range of the box operator \square . Here, J^{2m} denotes the truncation at degree $2m$ of the Taylor expansion at the origin. We then need to estimate U in terms of B . For that purpose, we shall first define a suitable analytic norm (depending on some “radius” r , see section 2.2) based on the scalar product we mentioned above (see section 2.1). Following (20), we shall then re-write the previous equation as

$$V = (\text{id} + Q_1 + Q_2)^{-1} \widetilde{\square}^{-1} B,$$

where $\widetilde{\square}$ denotes the restriction of the operator $\square := d_0 d_0^*$ to its image and Q_1, Q_2 are operators defined by (19) and (18), depending on f_m . Using **all properties of $\mathfrak{sl}(2, \mathbb{C})$ -triples** (see section 2.4), we shall be able to estimate the normal of Q_1, Q_2 and $\widetilde{\square}^{-1}$. As a consequence, we shall obtain an estimate of V in terms of B , and then an estimate of U in terms of B (see proposition 3.1). Then the conjugacy of V by $(\text{id} + U)^{-1}$ is normalized up to order $2m$: $(\text{id} + U)_*^{-1} V = NF_{2m} + R_{\geq 2m+1}$ where $NF_{2m} := N + f_{2m} N^*$ is a normal form. Choosing $R < r$ in an appropriate way, we shall show that the new normal form NF_{2m} as well as the new remainder $R_{\geq 2m+1}$ satisfy the same estimate w.r.t. the R -norm that NF_m and $R_{\geq m+1}$ satisfy w.r.t. the r -norm (proposition 4.1). This allow to do an induction process. The key point (lemma 4.2) is that the sequence of successive radii $\{R_k\}$ converges to a positive number. This is due to the sharp estimate of the solution U we obtain (for each step). A classical argument then shows that the sequence of (partial) normalizing diffeomorphism converges to a genuine holomorphic diffeomorphism that conjugates V to a normal form.

1.6 Convergence to the normal form requires a condition

We don't know whether condition (1) is necessary for the convergence of the normalizing transformation to hold (as it is the case of Bruno's condition (A) in the semi-simple case). However, Stróżyńska and Żoładek have shown in [SZ11], that the normalizing transformation of the vector field

$$(*) \begin{cases} \dot{x} &= y - x^2 + x^3 \\ \dot{y} &= z + 2x^4 \\ \dot{z} &= 0 \end{cases}$$

to its (generalized) Takens normal form is divergent at the origin. Although, we shall not investigate the dictionary between different “styles” of normal form, let us show that the “scalar product” normal form of (*) doesn’t satisfy (1). Indeed, we have the following

Lemma 1.5. *If $f(x, y, z)$ is a formal first integral of (*), then f is a function of z , only.*

Proof. Let \mathcal{L} be the vector field associated to (*) and let $N := y\frac{\partial}{\partial x} + z\frac{\partial}{\partial y}$ be its linear part at 0. Let us assume, by induction, that $f = \sum_{i=1}^k f_i z^i + \sum_{j \geq k+1} g_j$ where g_j is homogeneous of degree j in **all variables**, and f_i ’s are complex coefficients. Hence, we have $0 = \mathcal{L}(f) = \mathcal{L}(g_{k+1} + g_{k+2} + \text{h.o.t})$ so $N(g_{k+1}) = 0$. Therefore, g_{k+1} is a polynomial P_{k+1} of z and $h = xz - \frac{y^2}{2}$. Let us show that $P_{k+1} = \alpha_{k+1} z^{k+1}$ and that $g_{k+2} = \alpha_{k+2} z^{k+2} + \beta_{k+2} z^k h$. Indeed, the Taylor polynomial of degree $k+2$ of $\mathcal{L}(f)$ is

$$N(g_{k+2}) - x^2 \frac{\partial g_{k+1}}{\partial x} = 0 = N(g_{k+2}) - x^2 z \frac{\partial P_{k+1}}{\partial h}.$$

Setting $z = 0$, we obtain $y \frac{\partial g_{k+2}}{\partial x}(x, y, 0) = 0$. Therefore, $g_{k+2} = z \tilde{g}_{k+2}$ and $N(\tilde{g}_{k+2}) - x^2 \frac{\partial P_{k+1}}{\partial h} = 0$. Let us write $x^2 \frac{\partial P_{k+1}}{\partial h} = \alpha x^2 h^m + x^2 z p(z, h)$ for some polynomial p and a constant α . Since $x^2 h^m \in \text{Ker } N^*$, $x^2 \frac{\partial P_{k+1}}{\partial h}$ can belong to the range of N only when $\alpha = 0$. Thus, $x^2 \frac{\partial P_{k+1}}{\partial h} = x^2 z p(z, h)$, $\tilde{g}_{k+2} = zG$ and $N(G) = x^2 p(z, h)$. We continue this process. We finally obtain that $g_{k+2} = z^k G_2$ and $N(G_2) = \alpha x^2$ for some quadratic polynomial G_2 . Since $N^*(x^2) = 0$, we have $\alpha = 0$ and thus G_2 is a first integral : $G_2 = \alpha z^2 + \beta h$. This means that $g_{k+2} = \alpha_{k+2} z^{k+2} + \beta_{k+1} z^k h$. As a consequence, $N(g_{k+2}) = 0$, so that $\frac{\partial P_{k+1}}{\partial h} = 0$ and $P_{k+1} = \alpha_{k+1} z^{k+1}$. \square

As a consequence, system (*) can not be formally conjugate to a system satisfying (1). Indeed, if it was, then its formal first integral would be a push-forward of a first integral of the second system. Since (1) has $h = xz - \frac{y^2}{2}$ as first integral, then (*) would have a formal first integral of the form $h \circ \Phi$, for some formal diffeomorphism Φ , tangent to identity. But such a function cannot be a function of the variable z only.

Acknowledgement *Part of the work was done while the second author spent 6 months on a postdoctoral position provided by University of Nice Sophia Antipolis during academic year 2011-2012. The first author thank M. Gazor, J. Murdock and J. Sanders for interesting email exchanges on uniqueness of formal normal/hypernormal forms.*

2 Background

For any non-negative integer k , \mathcal{P}_k (resp. \mathcal{V}_k) denotes the space of homogeneous polynomials (resp. polynomial vector fields) of degree k . We shall also write $\mathcal{P}_{k,m} := \bigoplus_{j=k}^m \mathcal{P}_j$ (resp. $\mathcal{P}_{\geq k} := \bigoplus_{j=k}^{\infty} \mathcal{P}_j$) as well as $\mathcal{V}_{k,m} := \bigoplus_{j=k}^m \mathcal{V}_j$ (resp. $\mathcal{V}_{\geq k} := \bigoplus_{j=k}^{\infty} \mathcal{V}_j$)

2.1 Scalar product

We introduce first some notation. Let $\alpha \in \mathbb{N}^n$, we shall write $\alpha! := \alpha_1! \dots \alpha_n!$. We define \mathcal{P}_δ to be the space of homogeneous polynomials of degree δ in n variables x_1, \dots, x_n . We first consider the scalar product on \mathcal{P}_δ introduced by E. Fischer [Fis17] and G. Belitskii [Bel79] :

Lemma 2.1. *The scalar product defined on \mathcal{P}_δ :*

$$\left\langle \sum_{|\alpha|=\delta} a_\alpha x^\alpha, \sum_{|\beta|=\delta} c_\beta x^\beta \right\rangle_{B,\delta} := \sum_{|\alpha|=\delta} a_\alpha \bar{c}_\alpha \alpha!$$

has the following property : the operator $\frac{\partial}{\partial x_j}$ is adjoint to the multiplication by x_j operator, that is $\langle \frac{\partial f}{\partial x_j}, g \rangle_{B,\delta} = \langle f, x_j g \rangle_{B,\delta+1}$ and $\langle g, \frac{\partial f}{\partial x_j} \rangle_{B,\delta} = \langle x_j g, f \rangle_{B,\delta+1}$ for all $(f, g) \in \mathcal{P}_{\delta+1} \times \mathcal{P}_\delta$.

In [LS10], a variant of this scalar product was introduced, namely :

$$\left\langle \sum_{|\alpha|=\delta} a_\alpha x^\alpha, \sum_{|\beta|=\delta} c_\beta x^\beta \right\rangle_\delta = \sum_{|\alpha|=\delta} a_\alpha \bar{c}_\alpha \frac{\alpha!}{|\alpha|!}. \quad (3)$$

Let $\|\cdot\|_\delta$ be its associated norm. Since homogeneous polynomial of different degree are orthogonal to each other, we are able to define the norm of formal power series $f = \sum_{k \geq 0} f_k$ where f_k is a homogeneous polynomial of degree k as

$$\|f\|^2 := \sum_{k \geq 0} \|f_k\|_k^2.$$

It has the some important properties that are summarized in the following proposition :

Proposition 2.2. [LS10][proposition 3.6-3.7] *Let f, g be formal power series on \mathbb{C}^n . Let us write $f = \sum_{k \geq 0} f_k$ where f_k is a homogeneous polynomial of degree k . Then,*

- $\|fg\| \leq \|f\| \|g\|$
- f defines a germ of holomorphic function at the origin if and only if there exist positive constants M, C such that $\|f_k\|_k \leq MC^k$.

This scalar product induces a scalar product on \mathcal{V}_δ , the space of homogeneous vector fields of degree δ in n variables. Such a vector field V can be written as $V = \sum_{k=1}^n V_k \frac{\partial}{\partial x_k}$, where $V_k \in \mathcal{P}_\delta$.

$$\left\langle \sum_{k=1}^n V_k \frac{\partial}{\partial x_k}, \sum_{k=1}^n W_k \frac{\partial}{\partial x_k} \right\rangle_\delta = \sum_{k=1}^n \langle V_k, W_k \rangle_\delta. \quad (4)$$

2.2 The analytic norm

Let us define, for $\delta \in \mathbb{N}$, $c_\delta := \left(\sum_{|Q|=\delta} \frac{|Q|!}{Q!} \right)^{1/2}$. Let f be a formal power series vanishing at the origin. It can be written as a sum of homogeneous polynomials $f = \sum_{\delta \geq 1} f_\delta$, where $f_\delta \in \mathcal{P}_\delta$. We define, for $r \geq 0$,

$$|f|_r := \sum_{\delta \geq 1} \|f_\delta\|_\delta c_\delta r^\delta, \quad (5)$$

in this formula $\|f_\delta\|_\delta$ is the norm associated to the scalar product (3) :

$$\|f_\delta\|_\delta^2 = \left\| \sum_{|k|=\delta} f_k x^k \right\|_\delta^2 = \sum_{|k|=\delta} |f_k|^2 \frac{k!}{|k|!}.$$

In a similar manner, we extend this norm to the space of vector fields using (4).

Lemma 2.3. $c_\delta = n^{\delta/2}$.

Proof. This follows immediately by choosing $z_i = 1$, $1 \leq i \leq n$ in the formula (see, for instance, [IL05][lemma A.1.])

$$(z_1 + \dots + z_n)^\delta = \sum_{|Q|=\delta} \frac{|Q|!}{Q!} z^Q.$$

□

Lemma 2.4. *Let f, g be formal power series. Then $|fg|_r \leq |f|_r |g|_r$.*

Proof. We have

$$|fg|_r = \sum_{k \geq 0} \|\{fg\}_k\|_k c_k r^k = \sum_{k \geq 0} \left\| \sum_{i+j=k} f_i g_j \right\|_k c_k r^k.$$

We recall that $\{fg\}_k$ denotes the homogeneous part of degree k of fg in the Taylor expansion at the origin. Since the scalar product (3) defines a Banach algebra norm, we have $\left\| \sum_{i+j=k} f_i g_j \right\|_k \leq \sum_{i+j=k} \|f_i\|_i \|g_j\|_j$. Since $c_k = n^{k/2} = c_i c_j$ if $i+j=k$, then

$$|fg|_r \leq \sum_{k \geq 0} \sum_{i+j=k} \|f_i\|_i c_i r^i \|g_j\|_j c_j r^j = \left(\sum_{i \geq 0} \|f_i\|_i c_i r^i \right) \left(\sum_{j \geq 0} \|g_j\|_j c_j r^j \right)$$

□

Lemma 2.5. *Suppose that $|u|_R \leq r$, then $|f \circ u|_R \leq |f|_r$*

Proof. Let $f = \sum_{Q \in \mathbb{N}^n} f_Q x^Q$ be the Taylor expansion of f at the origin.

$$\begin{aligned}
|f \circ u|_R &= \left| \sum_{\delta \geq 1} \sum_{|Q|=\delta} f_Q \cdot (u(x))^Q \right|_R \leq \sum_{\delta \geq 1} \sum_{|Q|=\delta} |f_Q| \cdot |(u(x))^Q|_R \\
&\leq \sum_{\delta \geq 1} \sum_{|Q|=\delta} |f_Q| \cdot |(u(x))|_R^{|Q|} \leq \sum_{\delta \geq 1} \sum_{|Q|=\delta} |f_Q| r^{|Q|} \\
&\leq \sum_{\delta \geq 1} \sum_{|Q|=\delta} |f_Q| \sqrt{\frac{Q!}{|Q|!}} \sqrt{\frac{|Q|!}{Q!}} r^{|Q|} \\
&\quad \text{(Apply Cauchy Schwartz)} \\
&\leq \sum_{\delta \geq 1} \left(\sum_{|Q|=\delta} |f_Q|^2 \frac{Q!}{|Q|!} \right)^{1/2} \left(\sum_{|Q|=\delta} \frac{|Q|!}{Q!} r^{2|Q|} \right)^{1/2} \\
&\leq \sum_{\delta \geq 1} \left(\sum_{|Q|=\delta} |f_Q|^2 \frac{Q!}{|Q|!} \right)^{1/2} c_\delta r^\delta = \sum_{\delta \geq 1} \|f_\delta\|_\delta c_\delta r^\delta \\
&= |f|_r.
\end{aligned}$$

□

Lemma 2.6. *Let f be a polynomial of degree at most m , then $|\frac{\partial f}{\partial x_i}|_r \leq \frac{m}{r} |f|_r$.*

Proof. We remark first that it readily follows from Lemma 2.3 that $c_{\delta-1} \leq c_\delta$. We estimate

$$\begin{aligned}
\left| \frac{\partial f_\delta}{\partial x_j} \right|_{\delta-1}^2 &\leq \sum_{|Q|=\delta} |f_Q|^2 q_j^2 \frac{Q!}{q_j(\delta-1)!} \leq \sum_{|Q|=\delta} |f_Q|^2 \frac{Q!}{\delta!} \frac{\delta! q_j}{(\delta-1)!} \\
&\leq \sum_{|Q|=\delta} |f_Q|^2 \frac{Q!}{\delta!} (\delta q_j) \leq \delta^2 \sum_{|Q|=\delta} |f_Q|^2 \frac{Q!}{\delta!}.
\end{aligned}$$

Hence we have $|\frac{\partial f_\delta}{\partial x_i}|_{\delta-1} \leq \delta |f_\delta|_\delta$. As a consequence, we obtain the following estimates :

$$\begin{aligned}
\left| \frac{\partial f}{\partial x_i} \right|_r &= \sum_{\delta=1}^m \left| \frac{\partial f_\delta}{\partial x_i} \right|_{\delta-1} c_{\delta-1} r^{\delta-1} \leq \sum_{\delta=1}^m \delta |f_\delta|_\delta c_{\delta-1} r^{\delta-1} \\
&\leq \frac{m}{r} \sum_{\delta=1}^m |f_\delta|_\delta c_{\delta-1} r^\delta \leq \frac{m}{r} \sum_{\delta=1}^m |f_\delta|_\delta c_\delta r^\delta \\
&\leq \frac{m}{r} |f|_r.
\end{aligned}$$

□

2.3 Normal forms

Let $F = \sum_{i=1}^n \left(\sum_{j=1}^n f_{i,j} x_j \right) \frac{\partial}{\partial x_i}$ be a linear vector field on \mathbb{C}^n . It acts by derivation as a linear map from \mathcal{P}_δ to itself. According to lemma 2.1, the adjoint of this operator w.r.t. Belitskii scalar product is the derivation

$$F^* := \sum_{j=1}^n \left(\sum_{i=1}^n \bar{f}_{i,j} x_i \right) \frac{\partial}{\partial x_j}.$$

The adjoint w.r.t. the scalar product (3) is the same. Let \tilde{N} be a nilpotent linear vector field in \mathbb{C}^n . Let us first recall the fundamental result that links the $\mathfrak{sl}(2, \mathbb{C})$ -triple actions and adjoint operators with respect to the scalar product.

Proposition 2.1. *[BV12] Assume that \tilde{N} is regular. Then, there exists a linear change of coordinates L such that $N = L_* \tilde{N}$ and its adjoint N^* generate an $\mathfrak{sl}(2, \mathbb{C})$ -triple. More precisely, the differential operators $X = N^*$, $Y = N$ and $H = N^*N - NN^*$, acting on germs of holomorphic functions or formal power series, fulfill the $\mathfrak{sl}(2, \mathbb{C})$ -relations from formula (7). Furthermore, N acts on vector fields of $V \in \mathcal{V}_\delta$, $\delta \geq 2$, as*

$$d_{0,\delta}(V) := [N, V],$$

where $[\cdot, \cdot]$ denotes the Lie bracket of vector fields. Then, its adjoint $d_{0,\delta}^*$ w.r.t. to the scalar product (4) is $d_{0,\delta}^*(V) := [N^*, V]$ and the triple $X = d_{0,\delta}^*$, $Y = d_{0,\delta}$, $H = [d_{0,\delta}^*, d_{0,\delta}]$ satisfies the $\mathfrak{sl}(2, \mathbb{C})$ -relations from formula (7).

The operator d_0 is called the **cohomological operator** and we shall define the **box operator** of degree δ , to be

$$\square_\delta := d_{0,\delta} d_{0,\delta}^*. \quad (6)$$

We recall the main formal normal form result :

Proposition 2.2. *[LS10] Let $X := N + R_{\geq 2}$ be a nonlinear holomorphic (or formal) perturbation of the nilpotent linear vector field N in a neighborhood of the origin of \mathbb{C}^n . Then, there exists a formal diffeomorphism $\hat{\Phi}$ tangent to the identity that conjugates X to a formal normal form; that is $\hat{\Phi}_* X - N \in \text{Ker } \text{ad}_{N^*}$. Moreover, there exists a unique normalizing diffeomorphism $\Phi = \text{Id} + \mathcal{U}$ such that \mathcal{U} has a zero projection on the kernel of $d_0 = [S, \cdot]$.*

Definition 2.3. *The normal form of X obtained by conjugacy of the unique normalizing diffeomorphism $\Phi = \text{Id} + \mathcal{U}$ such that \mathcal{U} has a zero projection on the kernel of $d_0 = [S, \cdot]$, will be called **the normal form of X** .*

2.4 $\mathfrak{sl}(2, \mathbb{C})$ -representations

Throughout this section we work within the following setting. We consider a representation $\{X, Y, H\}$ of $\mathfrak{sl}(2, \mathbb{C})$ acting on a finite dimensional vector space \mathcal{V} , that is linear operators of \mathcal{V} and satisfy the relations

$$[X, Y] = H, \quad [H, X] = 2X, \quad [H, Y] = -2Y. \quad (7)$$

Such a family $\{X, Y, H\}$ is also called an $\mathfrak{sl}(2, \mathbb{C})$ -triple. The main definitions and propositions of this section can be found in [Bou90, Ser87].

Definition 2.7. *Let $\{X, Y, H\}$ be an $\mathfrak{sl}(2, \mathbb{C})$ -triple acting on a finite dimensional vector space \mathcal{V} . A nonzero vector b such that $Xb = 0$ and $Hb = \lambda b$ is called a primitive vector. The corresponding eigenvalue λ is called a weight.*

We use the following lemma from $\mathfrak{sl}(2, \mathbb{C})$ -representation theory:

Lemma 2.8. *[Ser87][IV-3] Let $\{X, Y, H\}$ be an $\mathfrak{sl}(2, \mathbb{C})$ -triple acting on a vector space \mathcal{V} . Let b be a primitive vector. Let us define*

$$e_m := \frac{Y^m b}{m!}, \quad v_m := Y^m b, \quad m \geq 0$$

and $e_{-1} := v_{-1} := 0$. The following properties hold:

- (a) $H(e_m) = (\lambda - 2m) e_m$, (a) $H(v_m) = (\lambda - 2m) v_m$,
- (b) $Y(e_m) = (m + 1) e_{m+1}$, (b) $Y(v_m) = v_{m+1}$,
- (c) $X(e_m) = (\lambda - m + 1) e_{m-1}$. (c) $X(v_m) = m(\lambda - m + 1) v_{m-1}$.

Proof. Let x be an arbitrary vector of \mathcal{V} such that $Hx = \lambda x$; then

$$HYx = [H, Y]x + YHx = -2Yx + \lambda Yx = (\lambda - 2)Yx.$$

As a consequence we obtain $He_m = (\lambda - 2m) e_m$ and $Y(e_m) = (m + 1) e_{m+1}$ follows from the definition of e_m . Finally we have that

$$\begin{aligned} mXe_m &= XYe_{m-1} = [X, Y]e_{m-1} + YXe_{m-1} \\ &= He_{m-1} + (\lambda - m + 2)Ye_{m-2} \\ &= m(\lambda - m + 1)e_{m-1}. \end{aligned}$$

The formulas for v_m are immediately deduced from the formulas for e_m . □

Corollary 2.9. *If \mathcal{V} is a finite dimensional vector space then the weight λ is an integer.*

Corollary 2.10. [Ser87][chap IV, theorem 2 and 3] Let e be a primitive vector and consider the vector subspace $W = \text{Span}\{e, Ye, Y^2e, \dots\}$ of \mathcal{V} ; then W is an irreducible $\mathfrak{sl}(2, \mathbb{C})$ -module. Every $\mathfrak{sl}(2, \mathbb{C})$ -module can be decomposed into a direct sum of irreducible $\mathfrak{sl}(2, \mathbb{C})$ -module.

Let us also assume that the vector space \mathcal{V} is provided with a scalar product $\langle \cdot, \cdot \rangle$ and an associated norm $\|\cdot\|$. Assume that the operators X, Y are adjoint w.r.t. the given scalar product : for each $v, w \in \mathcal{V}$, we have $\langle Xv, w \rangle = \langle v, Yw \rangle$, $\langle Yv, w \rangle = \langle v, Xw \rangle$.

Lemma 2.11. The operator $H = XY - YX$ is self-adjoint w.r.t. the scalar product.

Lemma 2.12. Let e_0, f_0 be two distinct primitive vectors of \mathcal{V} such that $\langle e_0, f_0 \rangle = 0$. Then, the following properties hold:

- (a) $\langle Y^m e_0, Y^n e_0 \rangle = 0$ for all $m, n \in \mathbb{N}$, $m \neq n$.
- (b) $\langle Y^m e_0, Y^n f_0 \rangle = 0$ for all $m, n \in \mathbb{N}$.

Proof. We show the first property by induction. We have

$$\langle Y e_0, e_0 \rangle = \langle e_0, X e_0 \rangle = 0$$

Let $n > m$, then it follows from Lemma 2.8 that

$$\langle Y^n e_0, Y^m e_0 \rangle = \langle Y^{n-1} e_0, (XY) Y^{m-1} e_0 \rangle = C_m \langle Y^{n-1} e_0, Y^{m-1} e_0 \rangle,$$

for a certain constant C_m that depends on m . Continuing this way concludes the first property. We proceed with the second property. Remark first that for all $n \geq 0$

$$\langle Y^n e_0, f_0 \rangle = \langle Y^{n-1} e_0, X f_0 \rangle = 0.$$

Let now $n \geq m$

$$\begin{aligned} \langle Y^n e_0, Y^m f_0 \rangle &= \langle Y^{n-1} e_0, (XY) Y^{m-1} f_0 \rangle \\ &= C_m \langle Y^{n-1} e_0, Y^{m-1} f_0 \rangle = \dots = C_m C_{m-1} \dots C_1 \langle e_0, Y^{m-n} f_0 \rangle = 0. \end{aligned}$$

□

Corollary 2.13. Let $\{X, Y, H\}$ be an $\mathfrak{sl}(2, \mathbb{C})$ -triple acting on a finite dimensional Hilbert space \mathcal{V} . Assume that the operators X, Y are adjoint w.r.t. the given scalar product. Then, there exists an orthogonal decomposition into irreducible $\mathfrak{sl}(2, \mathbb{C})$ -submodules of \mathcal{V} .

Proof. According to lemma 2.10, \mathcal{V} can be written as a direct sum of irreducible submodule W_i 's. Each of them is generated by a primitive element $b^{(i)}$ which is an eigenvector of H and belongs to the kernel of X . Let \tilde{V} be the vector space spanned by these $b^{(i)}$'s. Since $[H, X] = 2X$, the kernel of X is left invariant by H : if $Xf = 0$ then $-X(Hf) = 0$. In particular, \tilde{V} is invariant by H . The operator $H|_{\tilde{V}}$ is self-adjoint. Let $\{\tilde{b}^{(0)}, \dots, \tilde{b}^{(j)}\}$ be an orthonormal basis of eigenvectors of $H|_{\tilde{V}}$. Each $\tilde{b}^{(l)}$ is a primitive element and it generates an irreducible $\mathfrak{sl}(2, \mathbb{C})$ -module $\tilde{W}_l := \text{Span}\{Y^m(\tilde{b}^{(l)}) | m \in \mathbb{N}\}$. According to lemma 2.12, the \tilde{W}_l 's are pairwise orthogonal. Hence, $\mathcal{V} = \oplus W_i = \oplus^\perp \tilde{W}_i$. \square

We explain now how the norm acts on the individual irreducible representations $\text{Span}\{Y^l(b) | l \in \mathbb{N}\}$. We have the following lemma.

Lemma 2.14. *Suppose that b is a primitive element of \mathcal{V} and $Hb = \lambda b$ then for each $m \geq 0$ we have*

$$\|Y^m b\|^2 = \frac{m! \lambda!}{(\lambda - m)!} \|b\|^2. \quad (8)$$

Proof. According to lemma 2.8, we have that

$$\|Y^m b\|^2 = \langle Y^m b, Y^m b \rangle = \langle Y^{m-1} b, X Y^m b \rangle = m(\lambda - m + 1) \langle Y^{m-1} b, Y^{m-1} b \rangle.$$

It follows, by induction, that $\|Y^m b\|^2 = \frac{m! \lambda!}{(\lambda - m)!} \|b\|^2$. \square

Following the proof of lemma 2.13, let $\{b^{(0)}, \dots, b^{(j)}\}$ be a basis of orthogonal eigenvectors of the operator $H|_{\text{Ker}(X)}$. Let us define $b_m^{(i)} := Y^m(b^{(i)}) / \|Y^m(b^{(i)})\|$ for each $i \in \mathbb{N}$, $0 \leq i \leq j$. Using the same argumentation as in the proof of corollary 2.13 it follows that

$$\{b_m^{(i)} | m \in \mathbb{N}, 0 \leq i \leq j\} \setminus \{0\} \quad (9)$$

is a **basis of orthonormal eigenvectors of H** . We study the action of X , Y , H , XY and YX on this normed basis, as this will be convenient for later use.

Lemma 2.15. *Let $b \in \text{Ker}(X)$ be an eigenvector of H associated with eigenvalue λ . Let us set $b_m := Y^m(b) / \|Y^m(b)\|$, $m \geq 0$. Then the vectors b_m satisfy the following properties:*

$$\begin{aligned} H(b_m) &= (\lambda - 2m)b_m, \\ Y(b_m) &= \sqrt{(m+1)(\lambda - m)}b_{m+1}, \\ X(b_m) &= \sqrt{m(\lambda - m + 1)}b_{m-1}, \\ YX(b_m) &= m(\lambda - m + 1)b_m, \\ XY(b_m) &= (m+1)(\lambda - m)b_m. \end{aligned} \quad (10)$$

Proof. We use Lemma 8 and compute:

$$\begin{aligned}
H(b_m) &= \frac{H(Y^m(b))}{\|Y^m(b)\|} = (\lambda - 2m)b_m, \\
Y(b_m) &= \frac{Y(Y^m(b))}{\|Y^m(b)\|} = \frac{Y^{m+1}(b)}{\|Y^m(b)\|} = \frac{b_{m+1}\|Y^{m+1}(b)\|}{\|Y^m(b)\|} = \sqrt{(m+1)(\lambda-m)}b_{m+1}, \\
X(b_m) &= \frac{X(Y^m(b))}{\|Y^m(b)\|} = m(\lambda - m + 1)\frac{Y^{m-1}(b)}{\|Y^m(b)\|} \\
&= m(\lambda - m + 1)\frac{b_{m-1}\|Y^{m-1}(b)\|}{\|Y^m(b)\|} = \sqrt{m(\lambda - m + 1)}b_{m-1}, \\
YX(b_m) &= \sqrt{m(\lambda - m + 1)}Yb_{m-1} = m(\lambda - m + 1)b_m, \\
XY(b_m) &= \sqrt{(m+1)(\lambda - m)}Xb_{m+1} = (m+1)(\lambda - m)b_m.
\end{aligned}$$

□

It is a classical fact that XY is selfadjoint positive semi-definite such that $\mathcal{V} = \text{Ker}(XY) \oplus \text{Im}(XY) = \text{Ker}(Y)$, $\text{Im}(XY) = \text{Im}(X)$, $\text{Ker}(XY) = \text{Ker}(Y)$ (the same holds for YX if we interchange the role of X and Y).

3 The normal form procedure : The Newton method

We consider a germ of holomorphic vector field at the origin of \mathbb{C}^n , $V = N + R_{\geq 2}$, where N is its (nilpotent) linear part and $R_{\geq 2} = \sum_{k \geq 2} R_k$, with $R_k \in \mathcal{V}_k$, is a nonlinear perturbation of N . In this section, we investigate the conjugacy of V to a normal form. We assume that $\{N^*, N, D' := [N^*, N]\}$ (and, as a consequence $\{d_0^*, d_0, H' := [d_0^*, d_0]\}$) is an $\mathfrak{sl}(2, \mathbb{C})$ -triple acting on each space of homogeneous polynomials \mathcal{P}_k (resp. \mathcal{V}_k), $k \geq 2$. Hence, we can apply the entire abstract $\mathfrak{sl}(2, \mathbb{C})$ -theory developed in Section 2.4.

Let us assume that, at step m , we start with a germ of holomorphic vector field at the origin of the form $V_m = N + f_m N^* + R_{m+1}$. In this formula, $NF_m := N + f_m N^*$ is a polynomial normal form of degree m , $R_{m+1} := B + C$ where $B \in \mathcal{V}_{m+1, 2m}$, $C \in \mathcal{V}_{> 2m}$ and f_m is a polynomial of degree $m - 1$ vanishing at the origin and which is an invariant of both N and N^* , that is $N(f_m) = N^*(f_m) = 0$.

At each step m of the procedure, we try to find a suitable polynomial coordinate transformation $\Phi^{-1} = \text{id} + U$, where $U \in \mathcal{P}_{m+1, 2m}$ such that the diffeomorphism $\Phi^{-1} = \text{id} + U$ normalizes V_m up to order $2m$. This means that Φ^{-1} conjugates V_m to $V_{2m} := \Phi_* V_m = N + f_m N^* + \tilde{B} + \tilde{C}$ where $\tilde{B} \in \mathcal{V}_{m+1, 2m}$ is a normal form and $\tilde{C} \in \mathcal{V}_{> 2m}$. Let us set $R_{2m+1} := \tilde{C}$. The conjugacy equation reads :

$$(\text{id} + DU) \left(N + f_m N^* + \tilde{B} + \tilde{C} \right) = (N + f_m N^* + B + C) \circ (\text{id} + U).$$

The previous equation becomes :

$$\begin{aligned} NF_m + \tilde{B} + \tilde{C} + DU.(NF_m) + DU.(\tilde{B} + \tilde{C}) &= NF_m + D(NF_m).U + B \\ &+ (B \circ (\text{id} + U) - B) + C \circ (\text{id} + U) \\ &+ \{NF_m \circ (\text{id} + U) - NF_m - D(NF_m).U\}. \end{aligned} \quad (11)$$

On the other hand, we also have

$$\begin{aligned} \tilde{C} &= NF_m(\text{id} + U) - NF_m + (B + C) \circ (\text{id} + U) \\ &- \tilde{B} - DU.(NF_m + \tilde{B} + \tilde{C}). \end{aligned} \quad (12)$$

Hence,

$$\begin{aligned} \tilde{C} &= \int_0^1 D(NF_m).(y + tU(y))U(y) dt + (B + C) \circ (\text{id} + U) \\ &- \tilde{B} - DU.(NF_m + \tilde{B} + \tilde{C}). \end{aligned} \quad (13)$$

This will be used to estimate \tilde{C} within the Newton process. We rewrite equation (11) under the following form :

$$\begin{aligned} \tilde{B} - B + [NF_m, U] + \tilde{C} &= (B \circ (\text{id} + U) - B) + C \circ (\text{id} + U) \\ &+ \{NF_m \circ (\text{id} + U) - NF_m - DNF_m.U\} \\ &- DU(\tilde{B} + \tilde{C}). \end{aligned} \quad (14)$$

Since $U \in \mathcal{V}_{m+1,2m}$, the Taylor expansion at the origin of the right hand side of equation (14) contains only terms of order strictly larger than $2m$ at the origin. Therefore, we have

$$J^{2m}(\tilde{B} - B + [NF_m, U]) = 0,$$

where J^{2m} denotes the truncation at degree $2m$ of the Taylor expansion at the origin. We are led naturally to define the operator

$$\begin{aligned} d_{0,m+1,2m} : \mathcal{V}_{m+1,2m} &\rightarrow \mathcal{V}_{m+1,2m} \\ U &\mapsto J^{2m}([NF_m, U]). \end{aligned}$$

Let us set $Z := \pi_{\text{Im}(\square)}B$ and $U := d_0^*(V)$ for some $V \in \mathcal{V}_{m+1,2m}$. We want to solve the **iterated cohomological equation**

$$\pi_{\text{Im}(\square)}(J^{2m}([N + f_m N^*, d_0^*(V)])) = Z = \pi_{\text{Im}(\square)}J^{2m}(R_{m+1}). \quad (15)$$

The part that we do not remove from the conjugacy equation (14) is then given by

$$\tilde{B} := \pi_{\text{Ker}(\square)}B - \pi_{\text{Ker}(\square)}(J^{2m}([N + f_m N^*, d_0^*(V)])). \quad (16)$$

By assumption, we have that $\tilde{B} = g_{2m}N^*$ where $g_{2m} \in \mathcal{O}_N^N \cap \mathcal{O}_N^{N^*}$ is polynomial of degree $\leq 2m - 1$. We define $NF_{2m} := NF_m + \tilde{B}$.

The main result of this section is the estimate of the cohomological equation

Proposition 3.1. *There exists a positive constant η , independent of m such that if $1/2 \leq r \leq 1$ and $|D(NF^m - N)|_r + |NF^m - N|_r < \eta$, for any $Z \in \pi_{\text{Im}(\square)}(\mathcal{V}_{m+1,2m})$, then the unique solution $U = d_0^* V \in \mathcal{V}_{m+1,2m}$ of the “iterated cohomological equation” (15) :*

$$\pi_{\text{Im}(\square)}(d_{0,m+1,2m}(U)) = Z$$

satisfies

$$|U|_r \leq 2md|Z|_r. \quad (17)$$

The next section is devoted to the proof of this proposition but we start with some preparations. Recall that $\square = d_0 d_0^*$ is a self-adjoint, positive semi-definite operator and that we have the decomposition $\mathcal{P}_{m+1,2m} = \text{Im}(d_0) \oplus \text{Ker}(d_0^*) = \text{Im}(\square) \oplus \text{Ker}(\square)$ together with $\text{Im}(d_0) = \text{Im}(\square)$ and $\text{Ker}(d_0^*) = \text{Ker}(\square)$. As a consequence the operator

$$\tilde{\square} : \text{Im}(\square) \rightarrow \text{Im}(\square) : V \mapsto \square(V),$$

is an invertible operator. We define

$$\begin{aligned} P_1 : \mathcal{V}_{m+1,2m} &\rightarrow \mathcal{V}_{m+1,2m}; & P_2 : \mathcal{V}_{m+1,2m} &\rightarrow \mathcal{V}_{m+1,2m} \\ V &\mapsto J^{2m}(f_m d_0^* d_0^*(V)), & V &\mapsto J^{2m}(d_0^*(V)(f_m) N^*). \end{aligned} \quad (18)$$

We also define

$$\begin{aligned} Q_1 &:= \tilde{\square}^{-1} \pi_{\text{Im}(\square)} P_1|_{\text{Im}(\square)}, \\ Q_2 &:= \tilde{\square}^{-1} \pi_{\text{Im}(\square)} P_2|_{\text{Im}(\square)}. \end{aligned} \quad (19)$$

Hence, we have :

$$\begin{aligned} \pi_{\text{Im}(\square)} J^{2m}([N + f_m N^*, d_0^*(V)]) &= \pi_{\text{Im}(\square)} \left(\square(V) + J^{2m}(f_m d_0^* d_0^*(V)) - J^{2m}(d_0^*(V)(f_m) N^*) \right) \\ &= \square \pi_{\text{Im}(\square)} (\text{id} + Q_1 + Q_2)(V) \end{aligned} \quad (20)$$

We consider the operator $\text{id} + Q_1 + Q_2$ as acting on $\text{Im}(\square)$. If it is invertible then the solution of equation (15) is given by

$$V = (\text{id} + Q_1 + Q_2)^{-1} \tilde{\square}^{-1} Z. \quad (21)$$

We show in the next section that the operator $Q_1 + Q_2$ is, in fact, nilpotent and we give a bound on this operator. This allows to invert and to estimate the solution of the equation since it leads to:

$$(\text{id} + Q_1 + Q_2)^{-1} = \sum_{l=0}^{\alpha} (Q_1 + Q_2)^l.$$

Although, the degree of nilpotency α depends on m , **we essentially show, and this is the key step, that $(\text{id} + Q_1 + Q_2)^{-1}$ can be bounded independently of the degree m .** Such a sharp bound is needed to proof proposition 3.1.

3.1 Estimate of the solution of the iterated cohomological equation

This section is devoted to the proof of proposition 3.1.

We start by showing the nilpotency properties of several operators defined by (18), (19) acting on $\mathcal{V}_{m+1,2m}$.

Lemma 3.1. *The operators $P_1, P_2, Q_1, Q_2, P_1 + P_2, Q_1 + Q_2$ are nilpotent. Moreover P_1 and P_2 commute pairwise.*

Proof. Since f_m vanishes at the origin, the multiplication by f_m increases the order by 1. Therefore, the operator P_1 is nilpotent. The operator P_2 is nilpotent too. In fact, since $f_m \in \mathcal{O}_n^N \cap \mathcal{O}_n^{N^*}$, we have

$$\begin{aligned} P_2(P_2(V)) &= P_2(d_0^*(V)(f_m)N^*) = J^{2m}(d_0^*(d_0^*(V)(f_m)N^*)(f_m)N^*) \\ &= J^{2m}\left(N^*\left(d_0^*(V)(f_m)\right)N^*(f_m)N^*\right) = 0. \end{aligned}$$

To prove that P_1 and P_2 commute with each other is a little bit more intricate. We first compute

$$\begin{aligned} P_1P_2(V) &= J^{2m}(f_md_0^*d_0^*(d_0^*(V)(f_m)N^*)) = J^{2m}(f_mN^*N^*(d_0^*(V)(f_m))N^*) \\ &= J^{2m}(f_mN^*N^*N^*(V)(f_m)N^*). \end{aligned}$$

On the other hand, we have

$$\begin{aligned} P_2P_1(V) &= J^{2m}(d_0^*(f_md_0^*d_0^*(V))(f_m)N^*) = J^{2m}(f_md_0^*d_0^*d_0^*(V)(f_m)N^*) \\ &= J^{2m}(f_mN^*d_0^*d_0^*(V)(f_m)N^*) = J^{2m}(f_mN^*(N^*(N^*(V)(f_m)))N^*). \end{aligned}$$

This proves that P_1 and P_2 are pairwise commuting. Since P_1 is nilpotent there exists a natural number $N > 1$ (depending on m) such that $P_1^N = 0$. Since $P_2^2 = 0$ then $(P_1 + P_2)^{N+1} = \sum_{k=1}^{N+1} \binom{N+1}{k} P_1^k P_2^{N+1-k} = 0$.

Let us show that both P_1 and P_2 are upper triangular and \square is diagonal with respect to a well-chosen basis. To this end it suffices to write down the basis described in formula (9) ordered by increasing degree. Since the operators P_1 and P_2 increase the degree, they are automatically upper triangular in this basis. Furthermore it follows immediately from Lemma 2.15 that \square acts diagonally on this basis. One immediately deduces from these facts that the operators Q_1, Q_2 and $Q_1 + Q_2$ are upper triangular and hence nilpotent. \square

We recall that $Y = d_0, X = d_0^*, H' = [d_0^*, d_0] = -H$ defines an $\mathfrak{sl}(2, \mathbb{C})$ -triple acting on $\mathcal{V}_{m+,2m}$. Following corollary 2.13, we consider an orthonormal basis $\{b^{(1)}, \dots, b^{(k)}\}$ of primitive elements of $\mathcal{V}_{m+1,2m}$ (k depends on m) so that

$$\mathcal{V}_{m+1,2m} = \bigoplus_{1 \leq i \leq k}^{\perp} W_i$$

is an orthogonal decomposition into irreducible $\mathfrak{sl}(2, \mathbb{C})$ -submodules : we have, for $1 \leq i \leq k$,

$$W_i := \text{Span} \{b_n^{(i)}, n \in \mathbb{N}\} \quad \text{with} \quad b_n^{(i)} = \frac{d_0^n(b_0^{(i)})}{\|d_0^n(b_0^{(i)})\|}, n \in \mathbb{N}.$$

The set

$$\mathcal{B}_{m+1,2m} := \{b_n^{(i)} \mid 1 \leq i \leq k, n \in \mathbb{N}\} \setminus \{0\} \quad (22)$$

is an orthonormal basis of $\mathcal{V}_{m+1,2m}$ as immediately follows from Lemma 2.12.

The following lemma gives a bound for the operator Q_1 .

Lemma 3.2. *The operator Q_1 is bounded by*

$$6\|f_m\|. \quad (23)$$

Furthermore, we have

$$|Q_1(V)|_r \leq 6|f_m|_r|V|_r. \quad (24)$$

Proof. We recall the definition of the operator

$$Q_1(V) = \tilde{\square}^{-1} \pi_{\text{Im}(\square)} f_m d_0^* d_0^*|_{\text{Im}(\square)}(V).$$

Since $\square = d_0 d_0^*$ and since f_m is an invariant of both N and N^* , the multiplication operator by f_m and the operator \square commute pairwise. Therefore, we have

$$Q_1(V) = f_m \tilde{Q}_1(V), \quad \text{with} \quad \tilde{Q}_1(V) := \tilde{\square}^{-1} \pi_{\text{Im}(\square)} d_0^* d_0^*|_{\text{Im}(\square)}(V).$$

We “split” the operator \tilde{Q}_1 into two parts: $d_0^* d_0^*$ and $\tilde{\square}^{-1}$ and compute their actions using the decomposition into irreducible $\mathfrak{sl}(2, \mathbb{C})$ -modules. In particular, we compute their actions on the basis (22). We use the theory developed in Section 2.4 for that purpose.

According to lemma 2.15 and using the orthogonal basis $\mathcal{B}_{m+1,2m}$, we obtain for each j :

$$\begin{aligned} d_0^* d_0^* b_n^{(j)} &= X^2 b_n^{(j)} = \sqrt{(\lambda - n + 2)(n - 1)} \sqrt{(\lambda - n + 1)n} b_{n-2}^{(j)}. \\ \square b_n^{(j)} &= d_0 d_0^* = Y X b_n^{(j)} = n(\lambda - n + 1) b_n^{(j)}. \end{aligned}$$

Here, λ denotes the weight of the primitive element $b_0^{(j)}$. It follows that $Q_1(b_n^{(j)}) = 0$ for $0 \leq n \leq 2$ and for $n \geq 3$ that:

$$\tilde{Q}_1(b_n^{(j)}) = \frac{\sqrt{(\lambda - n + 2)(n - 1)} \sqrt{(\lambda - n + 1)n}}{(n - 2)(\lambda - n + 3)} b_{n-2}^{(j)}.$$

On the other hand, we have

$$\frac{\sqrt{(\lambda - n + 2)(n - 1)}\sqrt{(\lambda - n + 1)n}}{(n - 2)(\lambda - n + 3)} \leq 6.$$

Let $V = \sum_{j=1}^k \sum_{n \geq 0} V_n^j b_n^{(j)} \in \mathcal{V}_{m+1, 2m}^{(n)}$. According to the previous computations and estimates, we have

$$\left\| \tilde{Q}_1(V) \right\|^2 = \left\| \sum_{j=1}^k \sum_{n \geq 0} V_n^j \tilde{Q}_1(b_n^{(j)}) \right\|^2 \leq \sum_{j=1}^k \sum_{n \geq 0} 6^2 |V_n^j|^2 \|b_{n-2}^{(j)}\|^2 \leq 6^2 \|V\|^2.$$

Since \tilde{Q}_1 leaves each space of homogeneous vector fields invariant, we obtain as a consequence :

$$\begin{aligned} |Q_1(V_{m+1} + \dots + V_{2m})|_r &\leq |f_m|_r \left| \tilde{Q}_1(V_{m+1} + \dots + V_{2m}) \right|_r \\ &\leq |f_m|_r \sum_{k=m+1}^{2m} |\tilde{Q}_1(V_k)|_r = |f_m|_r \sum_{k=m+1}^{2m} \|\tilde{Q}_1(V_k)\|_k C_k r^k \\ &\leq 6|f_m|_r \sum_{k=m+1}^{2m} \|V_k\|_k C_k r^k = 6|f_m|_r |V|_r. \end{aligned}$$

This ends the proof of the lemma. \square

We proceed with an estimate on the operator Q_2 . Here the things tend to become a little more difficult since the operator Q_2 is not acting as a “ladder” operator as before. We can however make use of the following invariant subspace lemma :

Lemma 3.3. *The vector subspace of vector fields of the form $AN + BN^* + CH$, $A, B, C \in \mathcal{P}_{m+1, 2m}$, is invariant under the action of d_0, d_0^* .*

Proof. It is sufficient to show that this space is invariant by the action of d_0 and d_0^* . We recall that we consider the $\mathfrak{sl}(2, \mathbb{C})$ -triple $\{N^*, N, H'\}$, with $H' = -H = [N^*, N]$.

$$\begin{aligned} [N, AN + BN^* + CH'] &= N(A)N + N(B)N^* + N(C)H' + B[N, N^*] + C[N, H'] \\ &= (N(A) + 2C)N + N(B)N^* + (N(C) - B)H' \end{aligned} \quad (25)$$

$$\begin{aligned} [N^*, AN + BN^* + CH'] &= N^*(A)N + N^*(B)N^* + N^*(C)H' + A[N^*, N] + C[N^*, H'] \\ &= N^*(A)N + (N^*(B) - 2C)N^* + (N^*(C) + A)H'. \end{aligned} \quad (26)$$

\square

Lemma 3.4. *Let $||\nabla f_m|| = \left\| \frac{\partial f_m}{\partial x_1} \right\| + \dots + \left\| \frac{\partial f_m}{\partial x_n} \right\|$. The operator Q_2 is bounded by $C_0 ||\nabla f_m||$, for some positive constant C_0 , independent of m . Moreover, we have*

$$|Q_2(V)|_r \leq C_0 \left(\left| \frac{\partial f_m}{\partial x_1} \right|_r + \dots + \left| \frac{\partial f_m}{\partial x_n} \right|_r \right) \max(|N|_r, |N^*|_r, |H|_r) |V|_r$$

Proof. We recall the definition of the operator Q_2 .

$$Q_2(V) = \tilde{\square}^{-1} \pi_{\text{Im}(\square)} d_0^*(V) (f_m) N^*.$$

Lemma 3.3 above shows that the space of vector fields of the form $AN + BN^* + CH'$ is invariant under the action of the Lie algebra generated by d_0 and d_0^* . We compute the action of \square on this space and deduce the action of $\tilde{\square}^{-1}$ for elements of the form $B'N^* := d_0^*(V)(f_m)N^*$. Of course, we shall obtain conditions on B' in order to solve the equation. We then use this expression to compute a bound. Using equations (25) and (26), we deduce that

$$\begin{aligned} \square(AN + BN^* + CH') &= (NN^*(A) + 2(N^*(C) + A))N + (NN^*(B) - 2N(C))N^* \\ &\quad + (NN^*(C) + 2C + N(A) - N^*(B))H' \\ &=: A'N + B'N^* + C'H'. \end{aligned} \quad (27)$$

Let $D := NN^*$ be the operator acting on polynomials. Let us set $K := D + 2I$. In matrix notation, operator \square thus acts as

$$\begin{pmatrix} D + 2I & 0 & 2N^* \\ 0 & D & -2N \\ N & -N^* & D + 2I \end{pmatrix} \begin{pmatrix} A \\ B \\ C \end{pmatrix} = \begin{pmatrix} A' \\ B' \\ C' \end{pmatrix}. \quad (28)$$

We want compute the inverse of that operator for a vector of the type $B'N^* = [N^*, V](f_m)N^*$, so we set $A' = C' = 0$. Therefore, we have

$$KA = -2N^*(C). \quad (29)$$

Composing the third line by $Y = N$ and adding it to the second line leads to

$$N(D - 2NK^{-1}N^*)(C) = B'. \quad (30)$$

Finally, we have

$$D(B) = B' + 2N(C). \quad (31)$$

We first decompose V along the basis $\mathcal{B}_{m+1,2m}$ as defined by (22): $V = \sum_{i,n} V_n^{(i)} b_n^{(i)}$. Hence, we have $[N^*, V](f_m) = \sum_{i,n} V_n^{(i)} d_0^*(b_n^{(i)})(f_m)$. This motivates the following notation: For each irreducible submodule W_i of $\mathcal{V}_{m+1,2m}$ associated to the index $1 \leq i \leq k$, let λ be its weight (we omit to write i), we shall set for $0 \leq n \leq \lambda$:

$$a(n) := n(\lambda - n + 1).$$

We shall use the convention that $a(n) = 0$ and $b_n^{(i)} = 0$, for $n \leq 0$. Let us also set

$$v_n := d_0^*(b_n^{(i)})(f_m), \quad w_n := b_n^{(i)}(f_m).$$

Since f_m is an invariant of both N and N^* then

$$N(w_n) = [N, b_n^{(i)}](f_n) = \sqrt{(n+1)(\lambda-n)} b_{n+1}^{(i)}(f_m) = \sqrt{(n+1)(\lambda-n)} w_{n+1} \quad (32)$$

$$N^*(w_n) = [N^*, b_n^{(i)}](f_n) = \sqrt{n(\lambda-n+1)} b_{n-1}^{(i)}(f_m) = \sqrt{n(\lambda-n+1)} w_{n-1}. \quad (33)$$

$$\begin{aligned} H(w_n) &= N^*N - NN^*(w_n) = N^*(\sqrt{a(n+1)} w_{n+1}) - N(\sqrt{a(n)} w_{n-1}) \\ &= (\lambda - 2n) w_n. \end{aligned} \quad (34)$$

Since the elements w_n have distinct weights, those which are non-zero are linearly independent [Ser87][proposition 1, p.18]. Furthermore, we have

$$v_n = [N^*, b_n^{(i)}](f_m) = N^*(b_n^{(i)}(f_m)) = N^*(w_n) = \sqrt{a(n)} w_{n-1}.$$

Since we want to invert on elements which are orthogonal to the kernel of d_0^* , it is sufficient to consider $B'N^*$ as linear combination of element of the form $d_0^*(b_n^{(i)})(f_m)N^*$, $n \geq 2$. Indeed, we have $d_0^*(b_0^{(i)}) = 0$ (by definition) and $d_0^*(d_0^*(b_1^{(i)})(f_m)N^*) = 0$. Let us write

$$B' = \sum_i \sum_n B_n'^{(i)} b_n^{(i)}(f_m) = \sum_i \sum_n B_n'^{(i)} w_n.$$

Let us set $V = \sum_i \sum_{n \geq 1} V_n^{(i)} b_n^{(i)} \in \mathcal{V}_{m+1, 2m}$. Since $B' = [N^*, V](f_m)$ then, according to (33), we have

$$B_n'^{(i)} = V_{n+1}^{(i)} \sqrt{a(n+1)}. \quad (35)$$

We recall that $K = D + 2I$. The operators D and thus K are both diagonal when expressed in the basis $\{w_n\}$. Let us explain this in detail and let us compute their eigenvalues. Indeed, using Lemma 2.15 we have :

$$Kw_n = (NN^* + 2I)b_n^{(i)}(f_m) = [(d_0 d_0^* + 2I)b_n^{(i)}](f_m) = (a(n) + 2)w_n.$$

It follows that

$$K^{-1}w_n = \frac{w_n}{(a(n) + 2)}, \quad Dw_{n-1} = a(n-1)w_{n-1}.$$

As a consequence, we have

$$\begin{aligned} K^{-1}N^*w_{n-1} &= K^{-1}\sqrt{a(n-1)}w_{n-2} = \frac{\sqrt{a(n-1)}}{a(n-2)+2}w_{n-2}, \\ NK^{-1}N^*w_{n-1} &= \frac{\sqrt{a(n-1)}}{a(n-2)+2}Nw_{n-2} = \frac{a(n-1)}{a(n-2)+2}w_{n-1}, \\ (D - 2NK^{-1}N^*)w_{n-1} &= \frac{a(n-1)a(n-2)}{a(n-2)+2}w_{n-1}, \\ N(D - 2NK^{-1}N^*)w_{n-1} &= \frac{\sqrt{a(n)}a(n-1)a(n-2)}{a(n-2)+2}w_n. \end{aligned} \quad (36)$$

$$(37)$$

Therefore, according to (36), if

$$B_0^{(i)} = B_1^{(i)} = B_2^{(i)} = 0, \quad (38)$$

then equation (30) has a unique solution

$$C_{n-1}^{(i)} := \frac{a(n-2)+2}{\sqrt{a(n)}a(n-1)a(n-2)} B_n^{(i)}, \quad n \geq 3 \quad (39)$$

with $C_0^{(i)} = C_1^{(i)} = 0$. Since we have

$$K^{-1}N^*w_n = \frac{a(n)}{a(n-1)+2}w_{n-1},$$

then equation (29) has a unique solution given by

$$A_{n-1}^{(i)} := -2 \frac{a(n)}{a(n-1)+2} C_n^{(i)} = \frac{-2}{\sqrt{a(n+1)}a(n-1)} B_{n+1}^{(i)}, \quad n \geq 2 \quad (40)$$

and $A_0^{(i)} = 0$. Finally, we have for $n \geq 3$,

$$\begin{aligned} B_n^{(i)}w_n + 2C_{n-1}^{(i)}N(w_{n-1}) &= \left(B_n^{(i)} + 2\sqrt{a(n)}C_{n-1}^{(i)} \right) w_n \\ &= \left(1 + \frac{2(a(n-2)+2)}{a(n-1)a(n-2)} \right) B_n^{(i)}w_n, \\ B_n^{(i)}w_n + 2C_{n-1}^{(i)}N(w_{n-1}) &= 0, \quad n \leq 2, \\ D^{-1}(B_n^{(i)}w_n + 2C_{n-1}^{(i)}N(w_{n-1})) &= \frac{1}{a(n)} \left(1 + \frac{2(a(n-2)+2)}{a(n-1)a(n-2)} \right) B_n^{(i)}w_n, \quad n \geq 3. \end{aligned}$$

Therefore, the unique solution of (31) such that $B_0^{(i)} = B_1^{(i)} = B_2^{(i)} = 0$ is given by

$$B_n^{(i)} = \frac{1}{a(n)} \left(1 + \frac{2(a(n-2)+2)}{a(n-1)a(n-2)} \right) B_n^{(i)}, \quad n \geq 3. \quad (41)$$

Since λ is a positive integer, we have

$$\lambda \leq a(n) \leq \left(\frac{\lambda+1}{2} \right)^2, \quad 1 \leq n \leq \lambda.$$

Therefore, the numbers

$$\begin{aligned} \alpha_n &:= \frac{-2\sqrt{a(n+2)}}{\sqrt{a(n+1)}a(n-1)}, \quad n \geq 2 \\ \beta_n &:= \frac{\sqrt{a(n+1)}}{a(n)} \left(1 + \frac{2(1+\frac{2}{a(n-2)})}{a(n-1)} \right), \quad n \geq 3 \\ \gamma_n &:= \frac{\sqrt{a(n+1)}(1+\frac{2}{a(n-2)})}{\sqrt{a(n)}a(n-1)}, \quad n \geq 3, \end{aligned}$$

are uniformly bounded with respect to n and to λ . Hence, $\sup_n |\alpha_n|$, $\sup_n |\beta_n|$, $\sup_n |\gamma_n|$ are uniformly bounded with respect to m and i (the label of irreducible submodules of $\mathcal{V}_{m+1,2m}$). According to equations (35), (40), (41) and (39), we deduce that

$$\begin{aligned} A &= \sum_i \sum_{n \geq 2} \alpha_n V_{n+2}^{(i)} b_{n-1}^{(i)}(f_m) \\ B &= \sum_i \sum_{n \geq 3} \beta_n V_{n+1}^{(i)} b_n^{(i)}(f_m) \\ C &= \sum_i \sum_{n \geq 3} \gamma_n V_{n+1}^{(i)} b_{n-1}^{(i)}(f_m) \end{aligned}$$

solve equation (27) with $A' = C' = 0$ and $B' = d_0^*(V)(f_m)$ with restrictions (38).

Let $X = \sum_{j=1}^n X_j \frac{\partial}{\partial x_j}$ be a polynomial vector field. We have $\|X_j\| \leq \|X\|$ since $\|X\| = \sum_{j=1}^n \|X_j\|^2$. As a consequence, we have

$$\|X(f)\| = \left\| \sum_{j=1}^n X_j \frac{\partial f}{\partial x_j} \right\| \leq \sum_{j=1}^n \|X_j\| \left\| \frac{\partial f}{\partial x_j} \right\| \leq \|X\| \sum_{j=1}^n \left\| \frac{\partial f}{\partial x_j} \right\| \leq \|X\| \|\nabla f\|.$$

Let $V \in \mathcal{V}_{m+1,2m}$ be any unit vector field. It can be written along the orthonormal basis $\mathcal{B}_{m+1,2m}$ as $V := \sum_{(n,i)} V_{n,i} b_n^{(i)}$, where $\sum_{(n,i)} V_{n,i}^2 = 1$. We compute

$$\begin{aligned} \|Q_2(V)\| &\leq \left\| \sum_i \sum_{n \geq 2} V_{n+2,i} \alpha_n b_{n-1}^{(i)}(f_m) N \right\| + \left\| \sum_i \sum_{n \geq 3} V_{n+1,i} \beta_n b_n^{(i)}(f_m) N^* \right\| \\ &\quad + \left\| \sum_i \sum_{n \geq 3} V_{n+1,i} \delta_n b_{n-1}^{(i)}(f_m) H \right\| \\ &\leq \|\nabla f_m\| \cdot \|N\| \cdot \sup_n \alpha_n \|V\| + \|\nabla f_m\| \cdot \|N^*\| \cdot \sup_n \beta_n \|V\| \\ &\quad + \|\nabla f_m\| \cdot \|N^*\| \cdot \sup_n \delta_n \|V\| \\ &\leq C_0 \|\nabla f_m\|. \end{aligned}$$

Here C_0 is independent of m . We have hence shown that $\|Q_2\| \leq C_0 \|\nabla f_m\|$. Let us prove the second inequality. We first have

$$\left| \sum_{(n,i)} V_{n+2,i} \alpha_n b_{n-1}^{(i)}(f_m) N \right|_r \leq \left| \sum_{(n,i)} V_{n+2,i} \alpha_n b_{n-1}^{(i)}(f_m) \right|_r |N|_r.$$

Let us consider the polynomial vector field $X := \sum_{(n,i)} V_{n+2,i} \alpha_n b_{n-1}^{(i)}$. Let us show that

$$|X(f_m)|_r \leq |X|_r \left(\left| \frac{\partial f_m}{\partial x_1} \right|_r + \cdots + \left| \frac{\partial f_m}{\partial x_n} \right|_r \right).$$

Indeed, according to lemma 2.4, we have

$$|X(f_m)|_r \leq \max_j |X_j|_r \left(\left| \frac{\partial f_m}{\partial x_1} \right|_r + \cdots + \left| \frac{\partial f_m}{\partial x_n} \right|_r \right).$$

We have $|X_j|_r \leq |X|_r$ since $\|\{X_j\}_k\| \leq \|\{X\}_k\|$ where $\{X_j\}_k$ denotes the homogeneous component of degree k of X_j . As a consequence, we have

$$\begin{aligned} \left| \sum_{(n,i)} V_{n+2,i} \alpha_n b_{n-1}^{(i)}(f_m) N \right|_r &\leq \left| \sum_{(n,i)} V_{n+2,i} \alpha_n b_{n-1}^{(i)}(f_m) \right|_r |N|_r \\ &\leq \left| \sum_{(n,i)} V_{n+2,i} \alpha_n b_{n-1}^{(i)} \right|_r \left(\left| \frac{\partial f_m}{\partial x_1} \right|_r + \cdots + \left| \frac{\partial f_m}{\partial x_n} \right|_r \right) |N|_r. \end{aligned}$$

For $m+1 \leq k \leq 2m$, let I_k be the set of index i for which $b_n^{(i)}$ is homogeneous of degree k (for all n). Then, we have

$$\left| \sum_{(n,i)} V_{n+2,i} \alpha_n b_{n-1}^{(i)} \right|_r = \sum_{k=m+1}^{2m} \sum_{i \in I_k} \left\| \sum_n V_{n+2,i} \alpha_n b_{n-1}^{(i)} \right\| c_k r^k.$$

Since the $\mathcal{B}_{m+1,2m}$ is an orthonormal basis, we have

$$\left\| \sum_n V_{n+2,i} \alpha_n b_{n-1}^{(i)} \right\|^2 \leq \sum_n V_{n+2,i}^2 \alpha_n^2 \leq \max_n \alpha_n^2 \sum_n V_{n,i}^2.$$

Therefore, we have

$$\sum_{k=m+1}^{2m} \sum_{i \in I_k} \left\| \sum_n V_{n+2,i} \alpha_n b_{n-1}^{(i)} \right\| c_k r^k \leq \max_n \alpha_n \sum_{k=m+1}^{2m} \sum_{i \in I_k} \sqrt{\sum_n V_{n,i}^2} c_k r^k = \max_n \alpha_n |V|_r.$$

We obtain similar estimates for the other terms and we are done. \square

We recall the cohomological equation (21)

$$V = (\text{id} + Q_1 + Q_2)^{-1} \tilde{\square}^{-1} Z$$

where Z belongs to $\pi_{\text{Im}(\square)}(\mathcal{V}_{m+1,2m})$. According the previous lemmas, we have

$$|V|_r \leq \left[\sum_{k=0}^{\alpha(m)} (6|f_m|_r + C_0 |\nabla f_m|_r \max(|N|_r, |N^*|_r, |H|_r))^k \right] |\tilde{\square}^{-1} Z|_r \quad (42)$$

where we have written

$$|\nabla f|_r := \left(\left| \frac{\partial f_m}{\partial x_1} \right|_r + \cdots + \left| \frac{\partial f_m}{\partial x_n} \right|_r \right).$$

The aim of the remainder of the proof is to obtain an upper bound for the norm of the right hand side of (42). By definition, we have,

$$NF^m - N = f_m N^* = \sum_{i=1}^n f_m \left(\sum_{j=1}^n \tilde{s}_{i,j} x_j \right) \frac{\partial}{\partial x_i}$$

Let us set $L_i(x) := \sum_{j=1}^n \tilde{s}_{i,j} x_j$ for $1 \leq i \leq n$. We write f for f_m and we decompose $f = \sum_{k=1}^{m-1} f^{(k)}$ as a sum of homogeneous polynomial $f^{(k)}$ of degree k . We have

$$\begin{aligned} \|f L_i(x)\|^2 &= \langle L_i(x)f, L_i(x)f \rangle = \sum_{k=1}^m \langle L_i(x)f^{(k)}, L_i(x)f^{(k)} \rangle \\ &= \sum_{k=1}^m \frac{1}{(k+1)!} \left\langle f^{(k)}, \sum_{j=1}^n \tilde{s}_{i,j} \frac{\partial (L_i f^{(k)})}{\partial x_j} \right\rangle_B \\ &= \sum_{k=1}^m \frac{1}{(k+1)!} \left\langle f^{(k)}, \sum_{j=1}^n |\tilde{s}_{i,j}|^2 f^{(k)} + L_i(x) \sum_{j=1}^n \tilde{s}_{i,j} \frac{\partial f^{(k)}}{\partial x_j} \right\rangle_B. \end{aligned}$$

As a consequence, we have

$$\|f L_i(x)\|^2 = \sum_{k=1}^m \frac{1}{(k+1)!} \left(\left(\sum_{j=1}^n |\tilde{s}_{i,j}|^2 \right) \|f^{(k)}\|_B^2 + \|L_i^*(\partial) f^{(k)}\|_B^2 \right).$$

where $L_i^*(\partial)(f_k) = \sum_{j=1}^n \overline{\tilde{s}_{i,j}} \frac{\partial f_k}{\partial x_j}$. Hence, we have

$$\begin{aligned} |f L_i|_r &= \sum_{k=1}^{m-1} \langle L_i(x)f^{(k)}, L_i(x)f^{(k)} \rangle^{1/2} c_{k+1} r^{k+1} \\ &= \sum_{k=1}^{m-1} \left[\frac{1}{(k+1)} \left(\left(\sum_{j=1}^n |\tilde{s}_{i,j}|^2 \right) \|f^{(k)}\|^2 + \|L_i^*(\partial) f^{(k)}\|^2 \right) \right]^{1/2} c_{k+1} r^{k+1} \\ &\geq r \sum_{k=1}^{m-1} \frac{n^{1/2}}{\sqrt{k+1}} \left(\sum_{j=1}^N |\tilde{s}_{i,j}|^2 \right)^{1/2} \|f^{(k)}\| c_k r^k \\ &\geq r \left(\sum_{j=1}^n |\tilde{s}_{i,j}|^2 \right)^{1/2} |f|_r. \end{aligned}$$

Since $r \geq 1/2$ and since there exists an i for which $\sum_{j=1}^n |\tilde{s}_{i,j}|^2 \neq 0$, we finally obtain

$$|f_m|_r \leq \frac{2}{\sum_{j=1}^n |\tilde{s}_{i,j}|^2} |NF^m - N|_r. \quad (43)$$

We have $\frac{\partial f_m L_i}{\partial x_j} = f_m \frac{\partial L_i}{\partial x_j} + L_i(x) \frac{\partial f_m}{\partial x_j}$. According the previous estimate applied to $\frac{\partial f}{\partial x_j}$ instead of f , there exists an i such that, if $r \geq 1/2$, then

$$\left| \frac{\partial f_m}{\partial x_j} \right|_r \leq \frac{2}{\sum_{j=1}^n |\tilde{s}_{i,j}|^2} \left| \frac{\partial f_m L_i}{\partial x_j} - f_m \tilde{s}_{i,j} \right|_r.$$

As a consequence, there exists a positive constant c such that, for all $1 \leq j \leq n$ and all $m \geq 2$

$$\left| \frac{\partial f_m}{\partial x_j} \right|_r \leq c (|D(NF^m - N)|_r + |NF^m - N|_r) \quad (44)$$

Let us choose $\eta > 0$ such that

$$(6 + C_0 n c \max(|N|_1, |N^*|_1, |H|_1)) \eta < 1/2. \quad (45)$$

Using estimates (43) and (44) into (42), we obtain that if $1/2 \leq r \leq 1$ and $|D(NF^m - N)|_r + |NF^m - N|_r < \eta$, then

$$|V|_r \leq 2 |\tilde{\square}^{-1} Z|_r.$$

According to Lemma 2.6, since V is a polynomial vector field of degree $\leq 2m$ and $1/2 \leq r$, we have

$$|d_0^* V|_r \leq |V|_r |DN^*|_r + |DV|_r |N^*|_r \leq md |V|_r \quad (46)$$

for some positive constant d , independent of m . On the other hand, according to (10) we have :

$$|\tilde{\square}^{-1} Z|_r \leq |Z|_r$$

As a consequence, if $1/2 \leq r \leq 1$ and $|D(NF^m - N)|_r + |NF^m - N|_r < \eta$, then the unique solution $U = d_0^* V$ of the cohomological equation (15) satisfies

$$|U|_r \leq 2md |Z|_r \quad (47)$$

and we are done.

4 The iteration procedure towards convergence

We adapt the proof by the first author in [Sto00][section 8]. Let $1/2 < r \leq 1$ and $\eta > 0$ be a positive number that is small enough so that condition (45) is satisfied. For any integer $m \geq \lfloor 8n/\eta \rfloor + 1$. We define

$$\begin{aligned} \mathcal{NF}_m(r) &= \left\{ X \in \mathcal{V}_{>0} \mid \max(|X - N|_r, |D(X - N)|_r) < \eta - \frac{8n}{m} \right\}, \\ \mathcal{B}_{m+1}(r) &= \left\{ X \in \mathcal{V}_{>m} \mid |X|_r < 1 \right\}. \end{aligned}$$

Let $m = 2^k$ for some integer $k \geq 1$ and define

$$\rho = m^{-2/m}r \text{ and } R = \gamma_k m^{-4/m}r, \text{ where } \gamma_k = (2md)^{-1/m}.$$

Since $m^{1/m} \geq 1$ it is readily verified that $\rho < R < r \leq 1$ for m large enough, say $m \geq m_0$.

Let us go back to the framework of beginning of section 3. Suppose that we have already normalized our vector field up to order m . Our starting point is a vector field of the form $NF_m + R_{m+1}$, where $NF_m = N + f_m N^*$ is the polynomial part of degree m of the normal form and R_{m+1} is an analytic germ of order $\geq m+1$ at the origin. The following proposition will play the role of one step in the Newton process:

Proposition 4.1. *Assume that $NF_m \in \mathcal{NF}_m(r)$, $R_{m+1} \in \mathcal{B}_{m+1}(r)$. Suppose that m is large enough, say $m \geq m_0$, m_0 independent of r , then the unique $U \in d_0^*(\mathcal{V}_{m+1,2m})$ solution of (15) is such that:*

- (a) $\Phi := (id + U)^{-1}$ is a diffeomorphism such that $|id + U|_R < \rho$,
- (b) $\Phi_*(NF_m + R_{m+1}) = N_{2m} + R_{2m+1}$ is normalized up to order $2m$,
- (c) $NF_{2m} \in \mathcal{NF}_{2m}(R)$, $R_{2m+1} \in \mathcal{B}_{2m+1}(R)$.

Proof. According to the properties of the norm used (see section 2.2) and the definition of the new remainder (13), the proof of the first point, second point and of the inequality $R_{2m+1} \in \mathcal{B}_{2m+1}(R)$ of the proposition is identical to the proof of [Sto00][proposition 8.0.2]. Indeed, we obtained, from proposition 3.1, the estimate $|U|_r \leq \gamma_k^{-m}$.

We need to proof $NF_{2m} \in \mathcal{NF}_{2m}(R)$. Its proof slightly differs from the equivalent one in [Sto00][proposition 8.0.2]. We have to estimate the new normal form $NF^{2m} = NF^m + \tilde{B}$ as defined in (16). This expression contains a Lie bracket that we need to estimate. According to (16), we have

$$\tilde{B} = \pi_{\text{Ker}(\square)} B - \pi_{\text{Ker}(\square)} (J^{2m}([f_m N^*, d_0^*(V)])).$$

Indeed, since $\pi_{\text{Ker}(\square)}(d_0 d_0^*(V)) = 0$ (the range and kernel of \square are supplementary spaces), then $\pi_{\text{Ker}(\square)}(J^{2m}([N, d_0^*(V)])) = 0$. Hence, we have the following estimate :

$$|\tilde{B}|_R \leq |B|_R + |[f_m N^*, d_0^*(V)]|_R.$$

In order to estimate the right hand side, we write

$$[f_m N^*, d_0^*(V)] = f_m d_0^*(d^*(V)) - d_0^*(V)(f_m) N^*.$$

This leads to the following estimate :

$$|[f_m N^*, d_0^*(V)]|_R \leq |f_m|_R |d_0^*(d^*(V))|_R + |d_0^*(V)(f_m)|_R |N^*|_R.$$

According to (46) and (47), and since B (resp. f_m) has order $\geq m+1$ (resp. ≥ 1) at the origin, we have

$$\begin{aligned} |[f_m N^*, d_0^*(V)]|_R &\leq |f_m|_R 2(md)^2 |B|_R + 2md |B|_R |\nabla f_m|_R |N^*|_R \\ &\leq \left(\frac{R}{r}\right)^{m+2} (2(md)^2 |f_m|_r |B|_r + 2md |B|_r |\nabla f_m|_r |N^*|_r) \\ &\leq \left(\frac{1}{2md}\right)^{1+2/m} \frac{1}{m^{4(1+2/m)}} (2(md)^2 |f_m|_r |B|_r + 2md |B|_r |\nabla f_m|_r |N^*|_r) \end{aligned}$$

According to the assumption of proposition 3.1 and to the fact that $|B|_r \leq |B+C|_r \leq 1$, then for $1/2 \leq r \leq 1$, we have

$$|[f_m N^*, d_0^*(V)]|_R \leq C\eta \frac{1}{m^3}$$

for some positive constant C , independent of m . Furthermore, by lemma 2.6, we have

$$|D([f_m N^*, d_0^*(V)])|_R \leq C'\eta \frac{1}{m^2}$$

for some constant C' independent of m . This shows that $NF_{2m} \in \mathcal{NF}_{2m}(R)$ and we are done. \square

Let $1/2 < r \leq 1$ be a positive number and let us consider the sequence $\{R_k\}_{k \geq 0}$ of positive numbers defined by induction as follow :

$$\begin{aligned} R_0 &= r \\ R_{k+1} &= \gamma_k m^{-2/m} R_k \quad \text{where } m = 2^k \end{aligned}$$

Lemma 4.2. *The sequence $\{R_k\}_{k \geq 0}$ converges and there exists an integer m_1 such that for all integer $k > m_1$, $R_k > \frac{R_{m_1}}{2}$.*

Proof. We recall that $\gamma_k = (2md)^{-1/2^k}$. We have $R_{k+1} = r \prod_{i=1}^k \gamma_i (2^i)^{-2^{1-i}}$, thus

$$\ln R_{k+1} = - \sum_{i=1}^k \frac{\ln 2^{i+1} d}{2^i} - \ln c_1 \sum_{i=1}^k \frac{1}{2^i} - 2 \ln 2 \sum_{i=1}^k \frac{i}{2^i}$$

all the sums are convergent as k tends to infinity. It follows that there exists an integer m_1 , such that

$$\prod_{i=m_1+1}^{+\infty} \gamma_i (2^i)^{-2^{1-i}} > 1/2.$$

Thus, if $k > m_1$ then $R_k = R_{m_1} \prod_{i=m_1+1}^k \gamma_i (2^i)^{-2^{1-i}} > \frac{R_{m_1}}{2}$. \square

We are now in position to prove the theorem.

Let $N + R_{\geq 2}$ be a nonlinear holomorphic deformation of N in a neighborhood of the origin in \mathbb{C}^n . We may assume that it is holomorphic in a neighborhood of the closed polydisc D_1 . Let $m_2 = 2^{k_0}$ be the smallest power of 2 which is greater than $\max(m_0, 2^{m_1})$ where m_0 is the integer defined in Proposition 4.1. By a polynomial change of coordinates, we can normalize $N + R_{\geq 2}$ up to order m_2 : in these coordinates, $N + R_{\geq 2}$ can be written $NF^{m_2} + R_{m_2+1}$. If necessary, we may apply a diffeomorphism $a\text{Id}$ with $a \in \mathbb{C}^*$ sufficiently small so that $(NF^{m_2}, R_{m_2+1}) \in \mathcal{NF}_{m_2}(1) \times \mathcal{B}_{m_2+1}(1)$. We define as above the sequence $\{R_k\}_{k \geq k_0}$, with $R_{k_0} = 1$. Thus, for all integer $k > k_0$, we have $1/2 < R_k \leq 1$.

Then following [Sto00][p.202-203], we have that for all $k \geq k_0$, that there exists a diffeomorphism Ψ_k of $(\mathbb{C}^n, 0)$ such that $\Psi_k^*(NF^{m_2} + R_{m_2+1}) =: NF^{2^{k+1}} + R_{2^{k+1}+1}$ is normalized up to order 2^{k+1} , $(NF^{2^{k+1}}, R_{2^{k+1}+1}) \in \mathcal{NF}_{2^{k+1}}(R_{k+1}) \times \mathcal{B}_{2^{k+1}+1}(R_{k+1})$ and $|\text{Id} - \Psi_k^{-1}|_{R_{k+1}} \leq \sum_{p=k_0}^k \frac{1}{2^{2^p}}$.

Since $D(1/2) \subset D_{R_k}$ for all integers $k \geq k_0$, then the sequence $\{|\Psi_k^{-1}|_{1/2}\}_{k \geq k_0}$ is uniformly bounded. Moreover, the sequence $\{\Psi_k^{-1}\}_{k \geq k_0}$ converges coefficient wise to a formal diffeomorphism $\hat{\Psi}^{-1}$ (the inverse of the formal normalizing diffeomorphism). Therefore, this sequence converges in $\mathcal{H}_n^n(r)$ (for all $r < 1/2$) to $\hat{\Psi}^{-1}$ (see [GR71]). This means that the normalizing transformation is holomorphic in a neighborhood of $0 \in \mathbb{C}^n$ and the theorem has been proven.

References

- [Arn80] V.I. Arnold. *Chapitres supplémentaires de la théorie des équations différentielles ordinaires*. Mir, 1980.
- [Bel79] G. R. Belitskii. Invariant normal forms of formal series. *Funct. Anal. Appl.*, 13(1):46–47, 1979.
- [Bel82] G. R. Belitskii. *Normal forms relative to a filtering action of a group*, volume Transactions of the Moscow Mathematical Society, 1981, Issue 2, pages 1–39. American Mathematical Society, 1982. A translation of Trudy Moskov. Mat. Obšč. **40** (1979), Trans. Moscow Math. Soc. **1981**, no. 2 (1982).
- [Bou90] N. Bourbaki. *Groupes et Algèbre de Lie, Chapitres 7 et 8*. Masson, Paris, 1990.
- [Bru72] A.D. Bruno. Analytical form of differential equations. *Trans. Mosc. Math. Soc.*, 25,131-288(1971); 26,199-239(1972), 1971-1972.

- [BV12] P. Bonckaert and F. Verstringe. Normal forms with exponentially small remainder and gevreys normalization for vector fields with a nilpotent linear part. *Ann. Inst. Fourier (Grenoble)*, 62(6):2211–2225, 2012.
- [CDS04] M. Canalis-Durand and R. Schäfke. Divergence and summability of normal forms of systems of differential equations with nilpotent linear part. *Ann. Fac. Sci. Toulouse Math. (6)*, 13(4):493–513, 2004.
- [CG97] G. Cairns and E. Ghys. The local linearization problem for smooth $sl(n)$ -actions. *Enseignement Math.*, 43, 1997.
- [CS86] R. Cushman and J. A. Sanders. Nilpotent normal forms and representation theory of $sl(2, \mathbf{R})$. In *Multiparameter bifurcation theory (Arcata, Calif., 1985)*, volume 56 of *Contemp. Math.*, pages 31–51. Amer. Math. Soc., 1986.
- [Fis17] E. Fischer. Über die Differentiationsprozesse der Algebra. *J. für Math.* 148, 1-78., 148:1–78, 1917.
- [Gon95] X. Gong. Integrable analytic vector fields with a nilpotent linear part. *Ann. Inst. Fourier (Grenoble)*, 45(5):1449–1470, 1995.
- [GR71] H. Grauert and R. Remmert. *Analytische Stellenalgebren*. Springer-Verlag, 1971.
- [GS68] V.V. Guillemin and S. Sternberg. Remarks on a paper of Hermann. *Trans. Amer. Math. Soc.*, 130:110–116, 1968.
- [GY12] Majid Gazor and Pei Yu. Spectral sequences and parametric normal forms. *J. Differential Equations*, 252(2):1003–1031, 2012.
- [Her68] R. Hermann. The formal linearization of a semi-simple lie algebra of vector fields about a singular point. *Trans. Amer. Math. Soc.*, 130:105–109, 1968.
- [IL05] G. Iooss and E. Lombardi. Polynomial normal forms with exponentially small remainder for analytic vector fields. *J. Differential Equations*, 212(1):1–61, 2005.
- [Ito89] H. Ito. Convergence of Birkhoff normal forms for integrable systems. *Comment. Math. Helv.*, 64:412–461, 1989.
- [Ito09] H. Ito. Birkhoff normalization and superintegrability of Hamiltonian systems. *Ergodic Theory Dynam. Systems*, 29(6):1853–1880, 2009.
- [KOW96] H. Kokubu, H. Oka, and D. Wang. Linear grading function and further reduction of normal forms. *J. Differential Equations*, 132(2):293–318, 1996.

- [Kus67] A.G. Kushnirenko. Linear-equivalent action of a semi-simple lie group in the neighbourhood of a stationary point. *Funct. Anal. Appl.*, 1:89–90, 1967.
- [Lor06] F. Loray. A preparation theorem for codimension-one foliations. *Ann. of Math. (2)*, 163(2):709–722, 2006.
- [LS10] E. Lombardi and L. Stolovitch. Normal forms of analytic perturbations of quasihomogeneous vector fields: Rigidity, invariant analytic sets and exponentially small approximation. *Ann. Scient. Ec. Norm. Sup.*, pages 659–718, 2010.
- [Mar80] J. Martinet. Normalisation des champs de vecteurs holomorphes. *Séminaire Bourbaki*, 33(1980-1981),564, 1980.
- [MC88] R. Moussu and D. Cerveau. Groupes d’automorphismes de $(\mathbb{C}, 0)$ et équations différentielles $ydy + \dots = 0$. *Bull. Soc. math. France*, 16(1988),459-488, 1988.
- [Mos73] J. Moser. *Stable and random motions in dynamical systems*. Princeton University Press, Princeton, N. J.; University of Tokyo Press, Tokyo, 1973. With special emphasis on celestial mechanics, Hermann Weyl Lectures, the Institute for Advanced Study, Princeton, N. J, Annals of Mathematics Studies, No. 77.
- [Mur03] J. Murdock. *Normal forms and unfoldings for local dynamical systems*. Springer Monographs in Mathematics. Springer-Verlag, New York, 2003.
- [Mur04] James Murdock. Hypernormal form theory: foundations and algorithms. *J. Differential Equations*, 205(2):424–465, 2004.
- [Now94] A. Nowicki. *Polynomial derivations and their rings of constants*. Uniwersytet Mikołaja Kopernika, Toruń, 1994.
- [Rüs67] H. Rüssmann. Über die Normalform analytischer Hamiltonscher Differentialgleichungen in der Nähe einer Gleichgewichtslösung. *Math. Ann.*, 169:55–72, 1967.
- [San03] Jan A. Sanders. Normal form theory and spectral sequences. *J. Differential Equations*, 192(2):536–552, 2003.
- [Ser87] J.-P. Serre. *Complex Semisimple Lie Algebras*. Springer-Verlag, 1987.
- [Sie42] C.L. Siegel. Iterations of analytic functions. *Ann. Math.*, 43(1942)807-812, 1942.

- [SM71] C. L. Siegel and J. K. Moser. *Lectures on celestial mechanics*. Springer-Verlag, New York-Heidelberg, 1971. Translation by Charles I. Kalme, Die Grundlehren der mathematischen Wissenschaften, Band 187.
- [Sto00] L. Stolovitch. Singular complete integrability. *Publ. Math. I.H.E.S.*, 91:p.133–210, 2000.
- [Sto05] L. Stolovitch. Normalisation holomorphe d’algèbres de type Cartan de champs de vecteurs holomorphes singuliers. *Ann. of Math. (2)*, 161(2):589–612, 2005.
- [SŻ02] E. Stróżyńska and H. Żoładek. The analytic and formal normal form for the nilpotent singularity. *J. Differential Equations*, 179(2):479–537, 2002.
- [SŻ08] E. Stróżyńska and H. Żoładek. Multidimensional formal Takens normal form. *Bull. Belg. Math. Soc. Simon Stevin*, 15(5, Dynamics in perturbations):927–934, 2008.
- [SŻ11] E. Stróżyńska and H. Żoładek. Divergence of the reduction to the multidimensional nilpotent Takens normal form. *Nonlinearity*, 24(11):3129–3141, 2011.
- [Tak73] F. Takens. Singularities of vector fields. *I.H.E.S.*, 43(1973),47-100, 1973.
- [Vey78] J. Vey. Sur certains systèmes dynamiques séparables. *Am. Journal of Math.* 100, pages 591–614, 1978.
- [Vey79] J. Vey. Algèbres commutatives de champs de vecteurs isochores. *Bull. Soc. Math. France*, 107, pages 423–432, 1979.
- [Wei32] R. Weitzenböck. Über die Invarianten von linearen Gruppen. *Acta Math.*, 58(1):231–293, 1932.
- [Zun02] Nguyen Tien Zung. Convergence versus integrability in Poincaré-Dulac normal form. *Math. Res. Lett.*, 9(2-3):217–228, 2002.
- [Zun05] Nguyen Tien Zung. Convergence versus integrability in Birkhoff normal form. *Ann. of Math. (2)*, 161(1):141–156, 2005.